

Immunofluorescence Resource Guide

A Comprehensive Guide to Immunofluorescence Staining



YDLODEOHLQDQDGGDIURP

ELROQFDWHFKFKURPVSHFFRP

Making IF as easy as ABC

Helping you reach new visualization frontiers in your research: this is our mission. Since our founding in 1976, a primary driving principle has been to develop and manufacture labeling and detection technologies that make IF as easy as ABC.

- A.** Reliable and reproducible reagents that instill trust and confidence.
- B.** Simple and robust product designs that streamline workflows and allow elucidation of complex biological systems.
- C.** A knowledge base of over 100 years of combined IF experience to help you accelerate the pace of discovery.

It's as simple as that.

Vector Laboratories empowers scientific advances with innovative proteomic and glycomic solutions. Supporting scientific industries worldwide for 45 years and counting, Vector Laboratories is a trusted manufacturing partner with unmatched technical expertise and a culture of service. Customers rely on Vector Laboratories' immunohistochemistry, immunofluorescence, glycobiology, and bioconjugation products and custom manufacturing capabilities to move science forward with impact. Vector Laboratories' market-tested product portfolio provides the critical tools researchers need to precisely visualize and study tissues and cells as well as tackle today's biggest healthcare challenges. The company's products and technologies have been cited in more than 350,000 peer-reviewed publications, and its catalog and custom products are included in laboratory Standard Operating Procedures around the world. Learn more at vectorlabs.com.


Table of Contents

2	Immuofluorescence Introduction
2	Immuofluorescence Workflow
4	Immuofluorescence Selection Guide
6	Pioneering in IHC/IF Technology
8	Choosing a Detection System
9	Comparison of Detection Systems
10	Considerations for IF Detection
12	Fluorophore-Conjugated Secondary Antibodies
14	VectaFluor™ Ready-To-Use (R.T.U.) Antibody Reagents
	- VectaFluor R.T.U. Antibody Kits
	- VectaFluor Duet Immuofluorescence Double Labeling Kits
	- VectaFluor Excel Amplified Staining System
18	Fluorophore-Conjugated Streptavidin/Avidin Reagents
19	Anti-Streptavidin and Anti-Avidin Antibody Reagents
20	Secondary and Tertiary Detection Reagents
	- Biotinylated and Unconjugated Secondary Antibodies
	- Enzyme-Conjugated Secondary Antibodies
	- Avidin and Streptavidin Enzyme Conjugates
22	Species on Species Detection (Mouse)
23	M.O.M.® (Mouse on Mouse) Immunodetection Kits
24	Mounting Media
25	VECTASHIELD® Antifade Mounting Media
	- VECTASHIELD Antifade Mounting Medium
	- VECTASHIELD PLUS Antifade Mounting Medium
	- VECTASHIELD HardSet™ Antifade Mounting Medium
	- VECTASHIELD Vibrance® Antifade Mounting Medium
26	VECTASHIELD Mounting Media and Fluorophore Compatibility
27	VECTASHIELD Mounting Media Formats and Applications
28	VectaCell™ Trolox for Live Cell Imaging
	- VectaCell Trolox Antifade Reagent
29	Accessory Reagents
	- VECTABOND® Reagent Tissue Section Adhesive
	- ImmEdge® Hydrophobic Barrier Pen
	- ImmPrint™ Histology Pen
	- Control Antibodies
	- Antigen Unmasking Solutions
30	Blocking Background Signal
	- Vector® TrueVIEW® Autofluorescence Quenching Kits
	- BLOXALL® Endogenous HRP/AP Blocking Solution
	- Avidin/Biotin and Streptavidin/Biotin Blocking Reagents
	- Normal Sera and Bovine Serum Albumin (BSA)
	- Animal-Free Blocking Solutions
	- General Protein Blocking Reagents

Cover Image: Colon (FFPE): Antigen retrieved with Antigen Unmasking Solution (citrate-based, pH 6.0) and stained with CY5 Sambucus Nigra Lectin (SNA; fuchsia). VECTASHIELD Antifade Mounting Medium with DAPI counterstain (blue).

General Immunofluorescence Workflow

Vector Laboratories is your resource for premium labeling and detection products at each step of your IF workflow.



* For more information visit: vectorlabs.com/lectins

** TrueVIEW Autofluorescence Quenching is applied just prior to coverslipping

Secondary Antibody

- VectaFluor DyLight™ R.T.U. Secondary Antibodies
- VectaFluor Duet IF Double Labeling Kits
- M.O.M. (Mouse on Mouse) Immunodetection Kits
- Biotinylated secondary antibodies
- Fluorochrome-conjugated secondary antibodies
- Unconjugated secondary antibodies


Tertiary Reagent

- VectaFluor Excel DyLight Amplified Fluorescent Staining Systems
- Fluorochrome-conjugated avidin or streptavidin
- Biotinylated anti-avidin amplifying reagent
- Biotinylated anti-streptavidin amplifying reagent

Counterstain/ Mount

- VECTASHIELD Vibrance® Antifade Mounting Medium with or without DAPI counterstain (hard-setting)
- VECTASHIELD PLUS Antifade Mounting Medium with or without DAPI counterstain (non-hardening)
- VECTASHIELD Antifade Mounting Medium with or without DAPI counterstain (non-hardening)
- VECTASHIELD Antifade Mounting Medium with Propidium Iodide (PI) (non-hardening)
- VECTASHIELD HardSet Antifade Mounting Medium with or without DAPI counterstain (hard-setting)
- VECTASHIELD HardSet Antifade Mounting Medium with TRITC—Phalloidin (hard-setting)

Visualize



Immunofluorescence Selection Guide

Follow the simple steps below to choose the most appropriate labeling and detection solution for your experiment.

1

Choose Primary Antibody

- Specific for antigen of interest
- Consider tissue species and preparation (fixation)
- Consider antigen retrieval requirements



VECTABOND Reagent
(Tissue Section Adhesive)

Blocking Reagents




- Choices determined by the options selected in Steps 1-2
- Streptavidin/Biotin Blocking Kit (if using biotin/streptavidin system)
- Avidin/Biotin Blocking Kit
- Normal Sera (from the species of secondary antibody)
- M.O.M. (Mouse on Mouse) Blocking Reagent
- R.T.U. Animal-Free Blocker and Diluent
- BSA
- Casein Solution

SECONDARY DETECTION SYSTEMS

2

Choose Secondary Antibody and Tertiary Detection System

- Choose fluorophore based on wavelengths available in microscope
- Fluorophore-conjugated secondary antibody or biotinylated secondary antibody
- Consider sensitivity requirements
- Consider species of primary antibody
- Consider tissue species



One Step Single-label.

Fast. Convenient.

- Fluorophore-conjugated secondary antibodies
- VectaFluor R.T.U. DyLight Labeled Secondary Antibodies

or



One Step

Dual-label two-antigen detection.

Fast. Convenient.

- VectaFluor Duet IF Double Labeling Kits

or



Two Step

Biotin-based.

- Biotinylated secondary antibody and fluorophore conjugated avidin or streptavidin.
- See Step 3 for additional amplification

or



Two Step


Highest sensitivity. Non-biotin based.

- VectaFluor Excel Amplified Fluorescent Staining System (Amplifier Antibody + fluorescent tertiary antibody)

3

Choose Signal Amplification with Biotin-based Systems (Step 2, Option C)


- Multiple rounds of amplification possible (with biotin-based systems)



4

Reduce Autofluorescence from Aldehyde Fixation

- Vector TrueVIEW Autofluorescence Quenching Kits, with or without DAPI counterstain



5

Choose Mounting Media with or without a Counterstain


- VECTASHIELD Antifade Mounting Media, with or without counterstain



6

Visualize

- Fluorescence microscope
- View using appropriate excitation/emission filters




Tonsil (FFPE): Cytokeratin detected with R.T.U. VectaFluor Anti-Rabbit IgG, DyLight 488 (green) Kit. Autofluorescence quenched with TrueVIEW Quenching Kit and mounted with VECTASHIELD Vibrance Antifade Mounting Medium.

Legend



Primary antibody



Fluorophore-conjugated secondary antibody



Amplifier antibody



Biotinylated anti-avidin / streptavidin



Fluorophore



Biotin



Fluorophore-conjugated avidin / streptavidin



TrueVIEW Quenching Action

Pioneering in IHC/IF Technology

Observation is one of the fundamental steps in the scientific method. However, for centuries the scientific study of tissues was limited to observations of dissections with the unaided eye (gross anatomy).

This all changed in the 17th century when Anton Van Leeuwenhoek fabricated a microscope that allowed observations of tissues at the cellular level, thus establishing the science of histology. While early researchers found it relatively simple to distinguish between the cell boundaries and subcellular compartments in plants, doing so in animal tissue presented a much greater challenge. It wasn't until the late 19th century with the introduction of dyes, such as hematoxylin that Paul Mayer used to successfully stain nuclei, that the subcellular structure of tissues became visible and the science of histochemistry emerged.

The number of available tissue dyes and stains increased during the early 20th century, as did the number of molecular families they identified. However, the ability to identify individual cellular- or tissue-specific proteins remained elusive. This changed in the mid-20th century when Dr. Albert Coons demonstrated that fluorescently labeled antibodies could be used to localize bacteria inside macrophages, thus helping to pioneer the science of immunohistochemistry (IHC). Over the next two decades our understanding of antibodies, antigens and immunology grew rapidly. However, IHC remained largely a specialized research tool used primarily in university settings. Then in the late 1960's, Dr. Stratis Avrameas and Dr. Paul Nakane independently developed methods to covalently couple the enzyme horseradish peroxidase (HRP) to antibodies. HRP in the presence of diaminobenzidine and hydrogen peroxide creates a brown precipitate at the site of the HRP-conjugated antibody. The precipitate can be visualized using an ordinary light microscope. This allowed for the IHC results to be viewed in any lab having a light microscope, with no need for expensive, complicated fluorescence instrumentation.


The use of IHC as a research tool grew dramatically over the next decade. The technique began to be used in clinical settings at large university hospitals. The HRP assay system was further improved in the early 1980's when Dr. Su-Ming Hsu showed that the high affinity of avidin for biotin could be used to increase the stability of the enzyme antibody complex and improve the sensitivity of the assay. Vector Laboratories was instrumental in the development of the IHC field by commercializing such key technologies. The use of avidin- and biotin-based detection systems dominated the IHC market for the next two decades.

Up to this time, visualization using fluorescence microscopy was challenging due to the rapid photobleaching of fluorophores when exposed to the light of the microscope. This significantly limited the time over which a sample could be observed. In the early 1990's, VECTASHIELD Antifade Mounting Medium was introduced by Vector Laboratories as the first commercially available mountant for fluorescence. Not only did it have no autofluorescence (in the popular visualization channels), it was also effective in preventing the photobleaching, or fading of the fluorophores. This advancement in microscopy not only made image acquisition and analysis much more convenient, it provided researchers tools to challenge the limits of fluorescence detection.


In the last decade, immunofluorescence applications have been further improved by the adaptation of new super-resolution methods. Super-resolution microscopy allows imaging at a scale smaller than 200 nm. Due to its characteristics and convenience, VECTASHIELD Mounting Medium has been found to be quite suitable for super-resolution imaging methods like stochastic optical reconstruction microscopy (STORM) and structured illumination microscopy (3D-SIM). Olivier, et al., describes VECTASHIELD Mounting Medium as a "simple yet powerful buffer for 3D-STORM".

References


- Coons AH, Creech HJ, and Jones RN "Immunological properties of an antibody containing a fluorescent group" *Proc. Soc. Exp. Biol. Med.* 47, 200-202 (1941)
- Nakane P and Pierce GB Jr "Enzyme-labeled antibodies for the light and electron microscopic localization of tissue antigens" *J. Cell. Biol.* 33, 307-318 (1967)
- Leduc E, Avrameas S, and Bouteille M "Ultrastructural localization of antibody in differentiating plasma cells" *J. Exp. Med.* 127, 109-118. (1968)
- Hsu S-M, Raine L, and Fanger H "Use of Avidin-Biotin-Peroxidase Complex (ABC) in Immunoperoxidase Techniques: A Comparison between ABC and Unlabeled Antibody (PAP) Procedures" *J. Histochem. Cytochem.* 29(4), 577-580 (1981)
- Shi SR, Key ME, and Kalra KL "Antigen retrieval in formalin-fixed, paraffin-embedded tissues: an enhancement method for immunohistochemical staining based on microwave oven heating of tissue sections" *J Histochem Cytochem.* Jun, 39(6), 741-8 (1991)
- Bretschneider S, Eggeling C, and Hell SW "Breaking the Diffraction Barrier in Fluorescence Microscopy by Optical Shelving," *PRL* 98, 218103 (2007)
- Schermelleh L, Carlton PM, Haase S, Shao L, Winoto L, Kner P, Burke B, Cardoso MC, Agard DA, Gustafsson MGL, Leonhardt H, and Sedat JW "Subdiffraction multicolor imaging of the nuclear periphery with 3D structured illumination microscopy," *Science* 320, 1332-1336 (2008)
- Olivier N, Keller D, Rajan VS, Gönczy P, and Manley S "Simple buffers for 3D STORM microscopy," *Biochemical Optics Express* 4, 885-899 (2013)
- Wegel, E, et al. "Imaging cellular structures in super-resolution with SIM, STED and Localisation Microscopy: A practical comparison", *Scientific Reports*, 6, 27290 (2016)
- Childs GV "History of Immunohistochemistry" *Pathobiology of Human Disease* 3775-3796 (2014)




Fluorescent images with neon effect showing successive proliferation within the bulb of a hair follicle. Proliferating cells labeled for CldU (red), IdU (green) with cells dividing twice taking up both labels (yellow). Epidermal nuclei (blue) and dermal papilla nuclei (cyan) labeled with DAPI. Image provided by Nigel Hammond (Dixon Lab). Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK.



Skeletal muscle: Alpha-sarcoglycan (m), M.O.M. Fluorescein Kit (green) • Muscle-specific actin (m), M.O.M. Basic Kit, Texas Red™ Avidin DCS (red).



Rat muscle (FFPE): GFAP (red) and NF200 (green). Counterstained and coverslipped with VECTASHIELD Mounting Medium with DAPI (blue). The double IF was performed by Dr. Lynn Dong, Dept of Biomedical Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY, USA.



Astrocytes: stained for GFAP and detected with DyLight™ 488 labeled secondary antibody. mounted with VECTASHIELD HardSet Mounting Medium with DAPI. Image courtesy of Dr. Emma East, Dept. of Life Sciences, The Open University, U.K.



Choosing a Detection System

Immunofluorescence Overview

Immunofluorescence (IF) is a powerful method for visualizing proteins expressed directly within tissues. The IF method combines immunology and fluorescent molecules to localize proteins within defined morphological structures, and thus, provides insights into gene expression, protein-protein interactions, and biomarker identification. This method is used in a wide variety of applications, including basic research, assessment of normal and disease states in human and animal health, and in plant pathology studies.

Vector Laboratories develops and manufactures a wide selection of reagents for IF, including the novel TrueVIEW Autofluorescence Quenching Kit for removal of unwanted immunofluorescence. Also included are traditional fluorophore-conjugated antibodies and an extensive range of avidin/biotin products, which include conjugates with contemporary fluorophores such as DyLight dyes as well as kits that offer a significant increase in sensitivity or help streamline workflows. The VECTASHIELD, VECTASHIELD PLUS, VECTASHIELD HardSet and VECTASHIELD Vibrance Antifade Mounting Media are market-leading products on which researchers consistently rely to complete workflows and achieve signal retention for image acquisition and specimen archiving.


Comparison of Detection Systems

Choose the appropriate detection system for your experiment based on fluorophore (color), sensitivity, formats, flexibility, time and cost.


Detection System	Fluorophore	Color	Sensitivity	Concentrate	R.T.U. Format	Biotin-Free	Modular	Cost/ Assay
One-Step								
VectaFluor R.T.U. Secondary Antibodies	DyLight	Green Red	**		•	•		**
VectaFluor Duet IF Double Labeling Kits	DyLight	Green Red	**		•	•		***
Fluorophore-Conjugated Secondary Antibodies	Traditional† DyLight	Blue Green Orange Red Orange Red Far Red	**	•		•		•
Two-Step								
VectaFluor Excel Amplified kits	DyLight	Green Red	****		•	•		***
Streptavidin / Avidin Fluorophore Conjugates*	Traditional† DyLight Cyanine	Blue Green Red Far Red	***	•			•	•
M.O.M. (Mouse on Mouse) Immunodetection Kits								
M.O.M. (Mouse on Mouse) System	Traditional† DyLight	Blue Green Red Far Red	***	•			•	**

† Traditional fluorophores include: Fluorescein (FITC), Rhodamine (TRITC), Texas Red, AMCA, Phycoerythrin (PE)


* Sensitivity can be increased with multiple rounds of biotinylated anti-(strept)avidin and strept(avidin) fluorophore conjugates.



Fruit fly ovarian nurse cells. Cutaway three-dimensional reconstruction from a confocal stack of a *Drosophila melanogaster* ovary. E-cadherin (GFP, green), f-actin (red) and DAPI (blue). This image is a collaborative effort by Dr. Ian Newton and Dr. Paul Appleton, School of Life Sciences, University of Dundee, Dundee, UK.



Coronal section of a *Pdgfra/Rosa26* transgenic mouse brain at postnatal day (P15). The image depicts myelin basic protein (red) and oligodendrocyte precursor cells identified by the expression of GFP in the striatum. Cell nuclei are shown in blue. Image provided by Dr. Andrea Domenico Rivera, Institute of Biological and Biomedical Sciences, University of Portsmouth, Portsmouth, UK.



Adult organ of Corti labeled with anti-beta3-tubulin (green), phalloidin (cyan), anti-connexin 30 (red) and DAPI (blue). Image provided by Dr. Dan Jagger, Ear Institute, University College London, London, UK.

Considerations for IF Detection

Immunofluorescence detection reagents are used to localize and visualize target antigens expressed in tissue sections or cultured cells. When applied optimally, these highly specific reagents provide a defined contrast between their fluorescence, which demarcates the antigen, and the non-fluorescent region of the preparation. There are several options to achieve labeling for single and multiple antigen detection.

Direct Detection

One common IF method uses fluorophore-conjugated primary antibodies. This direct approach enables fast and easy IF visualization once the antibody has been conjugated; however, there are some disadvantages to this traditional method. For example, binding affinity and avidity could be compromised by the conjugation process, which would reduce signal and prevent moderately or weakly expressed antigens from being detected. Furthermore, expensive primary antibodies used at high concentrations could be cost prohibitive, and the visualization options would be limited to only one fluorophore.

Indirect Detection (One Step)


The indirect method, which uses labeled secondary antibodies, produces reliable, reproducible and economical IF results. This method avoids the disadvantages of directly conjugated primary antibodies and provides signal amplification that is necessary for most cell- and tissue-section labeling. Additionally, this one-step detection method is modular and allows simple substitution of the secondary with different fluorophore conjugates. Please refer to Table 2, page 13 for our range of concentrated reagents. Fluorophore-conjugated secondary antibodies would be recommended where a moderate to high expression of target antigen is expected.

Indirect Detection (Two Step)


Further signal amplification is introduced by using biotinylated secondary antibodies with avidin or streptavidin fluorophore conjugates. This well established and widely published methodology exploits the very high affinity between avidin or streptavidin and the small vitamin biotin. This two-step detection method enables the detection of weakly expressed antigens and provides a flexible and modular system with easy fluorophore substitution using different avidin or streptavidin conjugates (see pages 18-20). Additional amplification can be achieved by using biotinylated anti-avidin/streptavidin. For applications where use of biotin-based reagents for signal amplification would be problematic, we offer a non-biotin based two-step fluorescence approach with our VectaFluor Excel Amplified DyLight Antibody kits (see page 16).

Species Cross-Reactivity


Beyond the choices provided in the Selection Guide (pages 4-5), consideration should be given to the species of the tissue under examination and the species of the primary antibody. In cases of closely related species, it is recommended to use a secondary antibody that has been specifically adsorbed to remove cross-reacting antibodies. In instances where a mouse primary antibody is being applied to mouse tissue sections, it is recommended to use the M.O.M. Immunodetection System (see pages 22-23).



Staining of human breast cancer colony-forming culture for basal (Cytokeratin 14, green) and luminal markers (Cytokeratin 18, red). Image provided by Wendy Greenwood, method by Dr. Michael Prater, The Cancer Research UK Cambridge Institute, Cambridge, UK.



Hypoxia within hyperplastic breast tissue. A section of human breast tissue labeled for CAIX using immunofluorescence, counterstained and mounted using VECTASHIELD HardSet Antifade Mounting Medium with DAPI. Image provided by Dr. Carl Daly and Dr. Sarah Dean, Healthcare Science, University of the West of England, Bristol, UK.



Germinal center (GC) reaction in the spleen after acute viral infection. After recognition of viral antigens, T cells (blue) migrate from the T cell zone into the follicle where they interact with B cells (purple). The T cells 'help' B cells, instructing the formation of GCs (green) in which virus-specific B cells undergo selection, class switching and somatic hypermutation to secrete anti-viral antibodies to clear the infection. This work was conducted by Miriam Eckstein and Dr. Martin Vaeth, Department of Pathology, New York University, NY, USA.

Multiple Antigen Labeling

The visualization of two or more antigens on the same tissue section requires careful planning and specific reagent selection to generate unequivocal and reproducible staining results. We have recently introduced our VectaFluor Duet IF Double Labeling Kits that provide convenience and a straightforward approach to this often difficult and time-consuming application (see page 15).

Choosing fluorophores

Immunofluorescence detection reagents are labeled with fluorophores that absorb (excitation) and emit (emission) light at specific wavelengths. Fluorophores suitable for immunofluorescence are available across the complete visible light spectrum. The light source and filter cubes in a particular microscope must match the excitation and emission requirements of the specific fluorophore to achieve the optimal signal-to-noise ratios. For example, the absorption and emission peak wavelengths of fluorescein are 495 nm and 515 nm, respectively. Therefore, an excitation light source that is near 495 nm will yield the greatest emission signal. An emission filter that spans 515 nm will capture the emitted signal. These wavelengths are fixed properties of the fluorophores and the filter, and when properly paired, the system will yield the strongest signal and lowest background.

Table 1. Excitation and emission wavelengths and visual colors for immunofluorescence fluorophores.

Fluorophore	Color	Excitation Max (nm)	Emission Max (nm)
AMCA	Blue	350	450
DyLight 488	Green	493	518
Fluorescein	Green	495	515
CY3	Orange	550	570
Rhodamine	Orange	550	575
DyLight 549	Orange	562	576
Phycoerythrin	Red-Orange	565	574
DyLight 594	Red	593	618
Texas Red	Red	595	615
CY5	Far Red	649	670
DyLight 649	Far Red	652	672


Fluorophore-Conjugated Secondary Antibodies

All antibodies available from Vector Laboratories for immunological applications are prepared using optimized, proprietary immunization schedules that produce high-quality antibodies. The antibodies are affinity-purified, and solid-phase adsorption techniques are used to remove cross-reactivities that are likely to interfere with specific detection. The conjugated antibodies have the optimal degree of labeling to maximize signal output without compromising antibody specificity or affinity.


We offer researchers a range of traditional and contemporary conjugated fluorophores, including fluorescein, rhodamine, Texas Red, AMCA and phycoerythrin. DyLight dyes offer greater photostability, pH independence and brighter fluorescence than conventional fluorophores. DyLight dye-conjugated antibodies are ideal for cell- and tissue-based immunofluorescence and a variety of other applications. The DyLight dye conjugates are stable at pH 4-9 and compatible with many buffers and diluents.

Fluorophore-Conjugated Secondary (target species) Antibodies

- > Rabbit IgG
- > Mouse IgG
- > Human IgG
- > Goat IgG



Astrocytes: Stained for GFAP and detected with DyLight 594-conjugated secondary antibody. Mounted in VECTASHIELD HardSet Mounting Medium with DAPI. Image courtesy of Dr. Emma East, Department of Life Sciences, The Open University, Milton Keynes, UK.



Dorsal root ganglia cells (neurons and satellite glia): Beta III tubulin(ms), DyLight 549 Anti-Mouse IgG • s100(rb), DyLight 488 Anti-Rabbit IgG mounted in VECTASHIELD HardSet Mounting Medium with DAPI. Image courtesy of Dr. Emma East, Department of Life Sciences, The Open University, Milton Keynes, UK.

We offer a comprehensive range of fluorophore-conjugated secondary antibodies. These affinity-purified, highly specific antibodies, directed against the most commonly used primary antibody target species, are available with a wide choice of fluorophores and are presented in a concentrated format.

Table 2. Fluorophore-conjugated secondary antibodies.

Product	AMCA	Fluorescein	Texas Red™	DyLight™ 488	DyLight™ 549	DyLight™ 594	DyLight™ 649
Anti-Mouse IgG (H+L), made in horse		FI-2000	TI-2000	DI-2488	DI-2549	DI-2594	DI-2649
Anti-Rabbit IgG (H+L), made in horse				DI-1088		DI-1094	
Anti-Rabbit IgG (H+L), made in goat	CI-1000	FI-1000	TI-1000	DI-1488	DI-1549	DI-1594	DI-1649
Anti-Goat IgG (H+L), made in horse				DI-3088		DI-3094	
Anti-Goat IgG (H+L), made in rabbit	CI-5000						
Anti-Human Fluorophore-Conjugated Secondary Antibodies							
Anti-Human IgG (H+L), made in goat		FI-3000					
Anti-Human IgE, ε (Epsilon) chain specific, made in goat		FI-3040					
Anti-Human IgG, γ (Gamma) chain specific, made in goat		FI-3080					
Anti-Human κ (Kappa) Chain, made in goat	CI-3060						
Anti-Human Lambda Chain, made in goat	CI-3070	FI-3070					

VectaFluor™ Ready-To-Use Antibody Reagents


As investigators push research boundaries and require more sensitive, photo-stable fluorescent products, we have met this demand by developing a range of DyLight dye-conjugated secondary antibodies and novel detection kits that we have named VectaFluor reagents. The VectaFluor products are presented as pre-diluted, ready-to-use (R.T.U.) solutions that reduce optimization requirements at the researchers' end, thereby saving time and minimizing potential dilution errors which assists with greater consistency in collaborative efforts across lab environments.

Maximum performance is achieved when these VectaFluor reagents are used in combination with our VECTASHIELD Antifade Mounting Media (see pages 24-27).

VectaFluor R.T.U. Antibody Kits

The VectaFluor Ready-to-Use (R.T.U.) DyLight dye-conjugated secondary antibodies offer maximum convenience for fluorescence staining of cells and tissues. These affinity-purified, highly cross-adsorbed secondary antibodies are conjugated to DyLight dyes in a manner that ensures the maximum degree of labeling without compromising antibody affinity or specificity. DyLight dyes offer advantages such as bright fluorescence, excellent photostability and pH independence.

VectaFluor R.T.U. Antibody Reagents are suitable for use with rabbit, mouse, goat, sheep, and bovine IgG primary antibodies and are supplied as ready-to-use, pre-diluted, stabilized solutions (15 ml) with ready-to-use 2.5% normal horse serum (15 ml) for blocking.



Product	DyLight 488 (Green)	DyLight 594 (Red)
VectaFluor Anti-Rabbit IgG, made in horse	DI-1788	DI-1794
VectaFluor Anti-Mouse IgG, made in horse		DI-2794

VectaFluor R.T.U. Antibody Kits


- > Rabbit IgG
- > Mouse IgG
- > Goat IgG

DyLight 594 Kits

Excitation: 593 nm

Emission: 618 nm

Color: Red




Colon (FFPE): CD3 detected with R.T.U. VectaFluor Anti-Rabbit IgG, DyLight 594 (red) Kit. Mounted in VECTASHIELD HardSet Antifade Mounting Medium.

DyLight 488 Kits

Excitation: 493 nm

Emission: 518 nm

Color: Green



Tonsil (FFPE): Cytokeratin detected with R.T.U. VectaFluor Anti-Rabbit IgG, DyLight 488 (green) Kit. Autofluorescence quenched with TrueVIEW Quenching Kit and mounted with VECTASHIELD Vibrance Antifade Mounting Medium.

VectaFluor Duet Immunofluorescence Double Labeling Kits

Apply two colors in one step using the VectaFluor Duet IF Double Labeling Kits. These kits save time and effort in double-labeling immunofluorescence (IF) protocols, which can be long and tedious. The kits are configured to detect a mouse and a rabbit primary antibody with green and red fluorescence in one step.

- Two colors, one step
- Ready-to-use (R.T.U.)
- Robust cocktail formulation of DyLight anti-mouse IgG and DyLight anti-rabbit IgG


Two kit configurations are available:

Selection of a VectaFluor Duet IF Double Labeling Kit format is based on individual preference; however, certain parameters should be considered. For example, prevalence of the respective target antigens within a tissue section, and whether the more abundant antigen will be viewed by a green or red signal are important factors. Also consider possible overlap or co-localization of the antigens and which antibody combination would produce optimal results.

VectaFluor Duet Immunofluorescence Double Labeling Kit Contents:

- 15 mL 2.5% Normal Horse Serum for blocking, R.T.U.
- 15 mL VectaFluor Duet Reag-ent, R.T.U.


The affinity-purified, highly cross-adsorbed secondary antibodies are conjugated to DyLight dyes in a manner that maximizes the degree of labeling without compromising antibody affinity or specificity. The red and green DyLight dye-conjugated anti-mouse and anti-rabbit antibodies are then combined into a robust, stable cocktail formulation that yields sensitive and consistent dual staining. The VectaFluor Duet IF Double Labeling Kit is compatible with fluorescence staining of cells and tissues.




Product	Catalog Number
VectaFluor Duet IF Double Labeling Kit - DyLight 488 Anti-Rabbit (green) - DyLight 594 Anti-Mouse (red)	DK-8818
VectaFluor Duet IF Double Labeling Kit - DyLight 594 Anti-Rabbit (red) - DyLight 488 Anti-Mouse (green)	DK-8828

VectaFluor Duet Kits

- > Rabbit IgG (green)/
Mouse IgG (red)
- > Mouse IgG (green)/
Rabbit IgG (red)



Colon: Mouse Anti-Cytokeratin (AE1/AE3) and Rabbit Anti-Ki67 detected simultaneously with VectaFluor Duet IF Double Labeling Kit, DyLight 488 Anti-Mouse (green)/DyLight 594 Anti-Rabbit (red). Mounted in VECTASHIELD HardSet Mounting Medium.



Prostate: Rabbit Anti-PSA mAb and Mouse Anti-Smooth Muscle Actin detected simultaneously with VectaFluor Duet IF Double Labeling Kit, DyLight 488 Anti-Rabbit (green)/DyLight 594 Anti-Mouse (red). Mounted in VECTASHIELD HardSet Mounting Medium.

VectaFluor Excel Amplified Staining System


The VectaFluor Excel Amplified Staining System offers a convenient, non-biotin amplification method for fluorescence applications. This system uses an Amplifier Antibody - a specially prepared, high-affinity, unconjugated anti-mouse IgG or anti-rabbit IgG antibody produced in goat - followed by VectaFluor DyLight dye-conjugated anti-goat IgG antibody.

The affinity-purified, highly cross-adsorbed anti-goat IgG antibody is conjugated to DyLight dyes in a manner that ensures maximum degree of labeling without compromising antibody affinity or specificity. DyLight dyes offer advantages such as bright fluorescence, excellent photostability and pH independence.

- Stabilized, ready-to-use solutions
- Non-biotin signal amplification
- High sensitivity
- Low background

VectaFluor Excel Kit Contents:


- 15 ml 2.5% Normal Horse Serum for blocking, R.T.U.
- 15 ml Amplifier Antibody, R.T.U. (goat anti-mouse IgG or goat anti-rabbit IgG)
- 15 ml VectaFluor DyLight dye-conjugated Horse Anti-Goat IgG, R.T.U.




Product	DyLight 488 (Green)	DyLight 594 (Red)
VectaFluor Excel Amplified Anti-Rabbit IgG Kit	DK-1488	DK-1594
VectaFluor Excel Amplified Anti-Mouse IgG Kit	DK-2488	DK-2594

VectaFluor R.T.U Antibody Kits

- > Rabbit IgG (green or red)
- > Mouse IgG (green or red)



Tonsil: Anti-Multi-Cytokeratin, VectaFluor Excel Amplified DyLight 488 Anti-Mouse IgG Kit. Mounted in VECTASHIELD HardSet Mounting Medium with DAPI.



Tonsil: Anti-Multi-Cytokeratin, VectaFluor Excel Amplified DyLight 594 Anti-Mouse IgG Kit. Mounted in VECTASHIELD HardSet Mounting Medium with DAPI.

Frequently Asked Questions:

1) Can the VectaFluor Excel kits be applied to fixed cultured cells?

Yes, investigators have successfully applied these kits on fixed cultured cells directly, and cultured cells that have been formalin-fixed and paraffin-embedded. Please see references 1 and 2 below, respectively.

2) Are the VectaFluor Excel Kits compatible with other fluorescent secondary antibodies for double staining applications?

Yes, as indicated in reference 3 below. For this application to be successful however, investigators must use detection reagents raised in species that will not cross-react with the detection reagents of the VectaFluor Excel kit.

3) What are the advantages of using the VectaFluor Excel kits compared with secondary antibodies directly conjugated with fluorophores?

The main advantage of using the VectaFluor™ Excel kits is the increase in sensitivity the Amplifier Antibody generates. In most staining applications, investigators would see an increase of at least three- to four-fold over that of a secondary antibody directly conjugated with a fluorophore. This increase in sensitivity enables unambiguous visualization of weakly expressed antigens, as well as further dilution of potentially expensive primary antibodies.

4) Can the VectaFluor Excel kits be applied to any species of tissue?

The VectaFluor Excel kits were developed and optimized on human tissue sections. As with any secondary detection system, investigators should note potential cross-reactivity between the reagents being applied to a tissue section and inherent proteins. The VectaFluor Excel Anti-Rabbit IgG kits can be applied to rodent and primate species.

The VectaFluor Excel Anti-Mouse IgG kits are recommended for non-rodent tissues. Note however, that due to the VectaFluor Excel Anti-Goat IgG Reagent supplied in all VectaFluor Excel kits, recognition of proteins in goat, sheep, and bovine species may occur.

References:



- 1) Azumi E, et al. 2016. 75:97-104. *Orthodontic Waves*
- 2) Rengstl B, et al. 2017. 12(5): e0177378. *PLoS ONE*
- 3) Baillie R, et al. 2016. 69(8):742-744. *J. Clin. Pathol.*

Fluorophore-Conjugated Streptavidin/Avidin Reagents


The fluorophore-conjugated streptavidin and avidin reagents are highly purified and have low non-specific binding. These fluorescent conjugates can be used to detect biotinylated secondary antibodies and other macromolecules in various applications, including immunofluorescence, *in situ* hybridization and flow cytometry. The fluorescent signal can be amplified using biotinylated secondary antibodies and fluorophore-conjugated streptavidin or avidin.

Streptavidin/Avidin Fluorophores

- Blue (AMCA)
- Green (DyLight 488 and Fluorescein)
- Orange (DyLight 549 and CY3)
- Red (DyLight 594 and Texas Red)
- Far Red (DyLight 649, and CY5)



Tonsil (FFPE) was antigen-retrieved with Antigen Unmasking Solution and stained with Anti-Pan Cytokeratin (mouse; clone AE1/AE3), Biotinylated Horse Anti-Mouse IgG, and CY5 Streptavidin.



Tonsil (FFPE) was antigen retrieved with Antigen Unmasking Solution and stained with Anti-Pan Cytokeratin (mouse; clone AE1/AE3), Biotinylated Horse Anti-Mouse IgG and CY3 Streptavidin. Mounted in VECTASHIELD HardSet Mounting Medium with DAPI.


Product	AMCA	Fluorescein	Texas Red™	DyLight™ 488	DyLight™ 549	DyLight™ 594	DyLight™ 649	CY3	CY5
Streptavidin	SA-5008	SA-5001	SA-5006	SA-5488	SA-5549	SA-5594	SA-5649	SA-1300	SA-1500
Avidin		A-2001	A-2006						
Avidin DCS		A-2011	A-2016						

Anti-Streptavidin and Anti-Avidin Antibody Reagents


Use of the Biotinylated Anti-Streptavidin or Biotinylated Anti-Avidin antibodies is an ideal approach to increase sensitivity in (strept)avidin/biotin detection systems. These antibodies bind to streptavidin or avidin through both of their antigen-binding sites and the covalently-attached biotin residues. After the first application of a fluorophore-conjugated streptavidin or avidin, the signal is amplified by incubation with a Biotinylated Anti-Streptavidin or a Biotinylated Anti-Avidin antibody. That incubation is followed by a second incubation with fluorophore-conjugated streptavidin or avidin. This multi-layered approach accumulates more fluorophores at the target site and can provide a multi-fold amplification.

Biotinylated Anti-Avidin and Biotinylated Anti-Streptavidin amplification is ideal for the following applications:


- Immunofluorescence / Immunohistochemistry
- *In situ* hybridization
- Microarray assays
- ELISAs
- Blotting



Biotin-conjugated human chromosome 1 centromere-specific probe detected with Texas Red Avidin DCS, Biotinylated Anti-Avidin and Texas Red Avidin DCS (red). Mounted in VECTASHIELD Mounting Medium with DAPI (blue).



Biotin-conjugated human chromosome 1 centromere-specific probe detected with Fluorescein Avidin DCS, Biotinylated Anti-Avidin and Fluorescein Avidin DCS (yellow-green). Mounted in VECTASHIELD Mounting Medium with Propidium Iodide (red).



Product	Biotin	Unconjugated	DyLight™ 488
Anti-Streptavidin	BA-0500	SP-4000	SP-4488
Anti-Avidin	BA-0300		

Secondary and Tertiary Detection Reagents

Our secondary antibodies are prepared by hyper-immunizing animals in a manner that produces high affinity antibodies. These are then purified by an affinity chromatography procedure designed to remove any low-affinity antibodies. Cross-reactivities that can interfere with specific labeling are removed by solid-phase adsorption techniques. The final product is then subjected to rigorous quality-control assays including immunodiffusion, solid-phase enzyme immunoassays, gel electrophoresis, solid-phase binding assays and IHC tissue staining. These unconjugated antibodies are used to generate our enzyme conjugated and biotinylated secondary antibodies.

Biotinylated and Unconjugated Secondary Antibodies

Our high-affinity, purified, biotinylated and unconjugated secondary antibodies are manufactured under controlled conditions to retain maximum specificity and affinity. Our secondary antibodies are subjected to rigorous quality control assays and can be used for tissue and cell staining, ELISAs, and blotting.

Secondary Antibodies	Biotinylated					Unconjugated		
	Host Species (Concentrate)			Host Species (R.T.U.)†		Host Species (Concentrate)		
	Goat	Rabbit	Horse	Goat	Horse	Goat	Rabbit	Horse
Anti-Cat IgG (H+L)	BA-9000							
Anti-Chicken IgG (H+L)	BA-9010							
Anti-Goat IgG (H+L)		BA-5000	BA-9500		BP-9500		AI-5000	
Anti-Guinea Pig IgG (H+L)	BA-7000							
Anti-Hamster IgG (H+L)	BA-9100					AI-9100		
Anti-Horse IgG (H+L)	BA-8000							
Anti-Mouse IgG (H+L)	BA-9200		BA-2000	BP-9200	BP-2000	AI-9200		AI-2000
Anti-Mouse IgG (H+L), rat adsorbed			BA-2001					
Anti-Mouse IgM (H+L), Mu chain specific	BA-2020							
Anti-Rabbit IgG (H+L)	BA-1000		BA-1100	BP-9100	BP-1100	AI-1000		
Anti-Rat IgG (H+L)	BA-9400	BA-4000		BP-9400				
Anti-Rat IgG (H+L), mouse adsorbed	BA-9401	BA-4001					AI-4001	
Anti-Sheep IgG (H+L)		BA-6000						
Universal Anti-Mouse/Rabbit IgG (H+L)			BA-1400		BP-1400			
Universal Pan-Specific Anti-Mouse/Rabbit/Goat IgG (H+L)			BA-1300					

† Ready-to-use, prediluted stabilized solutions

Anti-Human Secondary Antibodies	Biotinylated	Unconjugated
	Host Species (Concentrate)	Host Species (Concentrate)
	Goat	Goat
Anti-Human IgG (H+L)	BA-3000	AI-3000
Anti-Human IgE, ε (Epsilon) chain specific	BA-3040	
Anti-Human IgG, γ (Gamma) chain specific	BA-3080	AI-3080
Anti-Human IgM, μ (Mu) chain specific	BA-3020	AI-3020
Anti-Human κ (Kappa) Chain, kappa chain specific	BA-3060	AI-3060

Enzyme-Conjugated Secondary Antibodies

Our high-affinity, purified antibodies are cross-linked with alkaline phosphatase (AP) or horseradish peroxidase (HRP) of the highest specificity. Our conjugation method ensures the maximum preservation of enzyme activity and antibody specificity. Recommended applications include tissue staining, ELISAs, and blotting.

Product	Catalog Number
Peroxidase	
Anti-Mouse IgG (H+L) made in horse Peroxidase-conjugated	PI-2000
Anti-Rabbit IgG (H+L) made in goat Peroxidase-conjugated	PI-1000
Anti-Human IgG (H+L) made in goat Peroxidase-conjugated	PI-3000
Anti-Goat IgG (H+L) made in horse Peroxidase-conjugated	PI-9500

Avidin and Streptavidin Enzyme Conjugates

Our enzyme-conjugated avidin and streptavidin are suitable for use in solid-phase assays, tissue- or cell-staining systems, and blotting. The conjugates are produced in optimized ratios with enzymes of the highest specific activity. Covalent linkages are specifically chosen to provide stable, highly active conjugates.

Product	Catalog Number
Alkaline Phosphatase	
Alkaline Phosphatase Streptavidin	SA-5100
Alkaline Phosphatase Avidin D	A-2100
Peroxidase	
Horseradish Peroxidase Streptavidin, concentrate	SA-5004
Horseradish Peroxidase Streptavidin, R.T.U.†	SA-5704
Horseradish Peroxidase Avidin D, concentrate	A-2004

† Ready-to-use, prediluted stabilized solutions

A microscopic image of tissue, likely a histological section, showing various cellular structures and patterns. A white rectangular text box is overlaid on the left side of the image, containing the title and introductory text. The background image is in grayscale and shows a complex network of cells and fibers.

Species on Species Detection (Mouse)


Solutions when your primary antibody is the same species as your specimen.

When a primary antibody is the same species as the specimen, the secondary antibody cannot distinguish between the endogenous immunoglobulins and the primary antibody. This can result in high background staining that obscures antigen-specific staining. Mouse on Mouse detection is especially important because of the vast number of primary antibodies made in mouse and the wide use of mice in model systems, xenografts, and other applications.


M.O.M. (Mouse on Mouse) Immunodetection Kits

The Vector M.O.M. Immunodetection kits are specifically designed to localize mouse primary antibodies on mouse tissue while avoiding background staining. These M.O.M. Kits contain our proprietary M.O.M. Mouse IgG Blocking Reagent. M.O.M. Kits are available based on either avidin-biotin technology (M.O.M. Elite Peroxidase Kit, Fluorescein Kit, or Basic Kit) or polymer technology (M.O.M. ImmPRESS® HRP Polymer Kit). Use the M.O.M. Immunodetection systems to introduce two or more different labels using a multiple antigen labeling protocol. You can detect several mouse primary antibodies on the same tissue section, regardless of the species of the tissue. Excellent staining results for a once difficult application have now become routine with the Vector Laboratories' M.O.M. System.

- Significantly reduces endogenous mouse Ig staining when using mouse primary antibodies on mouse tissue
- Simple protocols
- Eliminates tedious calculations
- Eliminates primary antibody prebinding steps
- Clear, crisp, specific staining of antigens of interest
- Compatible with fluorescent or enzyme-based detection
- Available with or without enzyme or fluorochrome





No M.O.M. Kit: Mouse Intestine (frozen) – Double label • Peripherin (m), Biotinylated Horse Anti-Mouse IgG, Fluorescein Avidin DCS (green) • Desmin (m), Biotinylated Horse Anti-Mouse IgG, Texas Red Avidin DCS (red). Note background and signal mixing.



With M.O.M. Kit: Mouse Intestine (frozen) – Double label • Peripherin (m), M.O.M. Fluorescein Kit (green) • Desmin (m), M.O.M. Basic Kit, Texas Red Avidin DCS (red). Compare with adjacent image prepared without M.O.M. Kit.

Recommended Applications

- Studies in genetically engineered mice
- Transgenic and knock-out models
- Mouse xenograft tissue
- Normal mouse tissue



Top: Mouse Colon: Multi-cytokeratin (m), M.O.M. Fluorescein Kit (green) • Desmin (m), M.O.M. Basic Kit, Texas Red Avidin DCS (red). Bottom: Mouse Tonsil: Ki67 (m), M.O.M. Fluorescein Kit (green) • CD20 (m), M.O.M. Basic Kit, Texas Red Avidin DCS (red). DCS (red).

Product	Catalog Number
M.O.M.® Elite® Immunodetection Kit, Peroxidase	PK-2200
M.O.M.® Immunodetection Kit, Fluorescein	FMK-2201
M.O.M.® Immunodetection Kit, Basic	BMK-2202
M.O.M.® ImmPRESS HRP Polymer Kit, Peroxidase	MP-2400
M.O.M.® Blocking Reagent	MKB-2213
M.O.M.® Biotinylated Anti-Mouse IgG Reagent*	MKB-2225
M.O.M.® ImmPRESS Polymer Reagent, Anti-Mouse IgG, Peroxidase	MPX-2402

* This reagent must be used with the M.O.M. Blocking Reagent (MKB-2213). It is not intended to be a stand-alone reagent for mouse on mouse applications.

A grayscale, high-magnification microscopic image of a cell, likely a neuron or similar cell type, showing a prominent nucleus with a nucleolus and surrounding cytoplasm with various organelles. The image is used as a background for the text.

Mounting Media

Choosing an effective mounting medium is especially important for immunofluorescence imaging. Fluorophores are susceptible to photobleaching and fading from both the imaging excitation light and during storage. The right mounting medium will protect your samples for short- and long-term use and archiving.

VECTASHIELD Antifade Mounting Media

VECTASHIELD Antifade Mounting Media formulations offer unsurpassed protection against fading and photobleaching. The VECTASHIELD, VECTASHIELD PLUS, VECTASHIELD HardSet and VECTASHIELD Vibrance Antifade Mounting Media are well-established, market-leading products that complete the workflow and provide excellent signal retention for image acquisition and specimen archiving.

- Inhibits photobleaching of most fluorophores, dyes, fluorescent proteins and stains
- Ideal refractive index
- Ready-to-use, no warming necessary
- Continues to inhibit photobleaching even after prolonged storage
- With or without nuclear or cytoskeletal counterstain
- Hardening or non-hardening formulations and easy-to-use

VECTASHIELD Antifade Mounting Medium

VECTASHIELD Antifade Mounting Medium is a glycerol-based, aqueous mountant that remains a viscous liquid on the slide rather than solidifying. After mounting, cover-slipped slides will not readily dry out, enabling you to review them for weeks without the need for sealing. For prolonged storage, coverslips can be permanently sealed with nail polish applied on the coverslip perimeter.

VECTASHIELD PLUS Antifade Mounting Medium


VECTASHIELD PLUS is a new formulation of non-setting media that improves upon the original VECTASHIELD products. Specifically, VECTASHIELD PLUS exhibits no inherent background or toning and provides superior fluorophore signal retention across the spectrum, including far-red wavelengths.

VECTASHIELD HardSet Antifade Mounting Medium


VECTASHIELD HardSet Antifade Mounting Medium is an aqueous mountant that hardens at room temperature in as little as 20 minutes. This mounting medium provides easy slide handling, eliminates the need to secure the coverslip with nail polish, and is convenient for use with oil immersion microscopy. Available with or without DAPI or TRITC-phalloidin counterstain.

VECTASHIELD Vibrance Antifade Mounting Medium


VECTASHIELD Vibrance Antifade Mounting Medium is an aqueous mountant that hardens at room temperature in as little as 20 minutes. This mounting medium provides easy slide handling, eliminates the need to secure the coverslip with nail polish, and is convenient for use with oil immersion microscopy. Available with or without DAPI counterstain.



Human Colon: Rabbit Anti-Cytokeratin (AE1/AE3) and Mouse Anti-Desmin detected simultaneously with VectaFluor Duet Double Labeling Kit; Vasculature detected using DyLight 649 UEA I Lectin (purple). Mounted in VECTASHIELD PLUS Antifade Mounting Medium.



Mouse embryonal fibroblasts: Anti-Integrin (m) detected with DyLight 488 Anti-Mouse IgG, mounted in a 1:1 mixture of VECTASHIELD HardSet Mounting Medium with DAPI and VECTASHIELD HardSet Mounting Medium with TRITC-Phalloidin.




Human uterine section (FFPE): Stained for desmin (green) and dytokeratin (red) using VectaFluor Duet Double Labeling Kit, and vasculature using DyLight 649 UEA I lectin (purple). Mounted in VECTASHIELD Vibrance Antifade Mounting Medium with DAPI (blue).

Product	No Counterstain	DAPI	PI	TRITC-Phalloidin
VECTASHIELD® Antifade Mounting Medium (non-hardening)	H-1000	H-1200	H-1300	
VECTASHIELD® PLUS Antifade Mounting Medium (non-hardening)	H-1900	H-2000		
VECTASHIELD® HardSet™ Antifade Mounting Medium (hardening)	H-1400	H-1500		H-1600
VECTASHIELD Vibrance® Antifade Mounting Medium (hardening)	H-1700	H-1800		

VECTASHIELD Mounting Media and Fluorophore Compatibility

VECTASHIELD Mounting Media are the most widely referenced antifade mounting media for immunofluorescence applications. Currently over 60,000 published references cite VECTASHIELD Mounting Media and describe compatibility with over 130 fluorophores and fluorescent markers. This data underscores the prominence of VECTASHIELD reagents in this application.


The graphic below highlights the most commonly referenced fluorophores used in combination with VECTASHIELD Antifade Mounting Media.



The fluorescent compounds listed in the table below are select reagents that are also cited as being successfully used in combination with VECTASHIELD Antifade Mounting Media. The range of these compounds, from traditional to contemporary, across a broad spectral range, and used in an array of applications, showcase the versatility of VECTASHIELD reagents. Over 130 fluorophores and fluorescent markers have been used with VECTASHIELD products please see a the complete [Compatibility List](#) available on our website.

Fluorophore				
acridine orange	coumarin	Fluoro-Jade®	NeuroTrace®	Quantum dot/Qdot
Alexa Fluor 350	dihydroethidium	Lucifer yellow	Nile red	SYTOX® Green
Alexa Fluor 680	DRAQ5™	LysoTracker®	Oil red O	TAMRA
Atto® dyes	Evans blue	LysoTracker® Red	Pacific Blue™	thioflavin s
BODIPY®	fast blue	MitoTracker® Red	PicoGreen®	TOTO®-3

VECTASHIELD Mounting Media Formats and Applications



The illustration above features established applications for our antifade mounting media formats. VECTASHIELD Antifade Mounting Media are widely utilized to protect the inherent fluorescent properties of traditional and contemporary fluorophores in many applications using epifluorescence and confocal microscopy.

The versatility of the original VECTASHIELD format solves the demands of labs and core facilities using multiple platforms and fluorescent markers. Furthermore, VECTASHIELD reagents are also recognized as leading media in emerging techniques such as super resolution microscopy (SRM).

Of the SRM techniques currently being performed, the properties of VECTASHIELD Antifade Mounting Media have been found to be advantageous in stochastic optical reconstruction microscopy (STORM) and structured illumination microscopy (SIM).

* Super Resolution (STORM and SIM) select references:

Olivier N, Keller D, Rajan VS, Gönczy P, and Manley S. 2013. "Simple buffers for 3D STORM microscopy." 4, 885-899. *Biochemical Optics Express*.

Wegel, E., et al. 2016. "Imaging cellular structures in super-resolution with SIM, STED and Localisation Microscopy: A practical comparison", 6, 27290. *Scientific Reports*.

VectaCell Trolox for Live Cell Imaging

Whereas immunofluorescence staining gives a snapshot of a cell or tissue at a specific time point, live cell imaging allows the observation of biological processes over a period of time. This is important for studying biological functions, interactions, and structures in various applications (e.g., the effects of drugs and other biomolecules).

VectaCell Trolox enables and enhances live cell imaging studies. VectaCell Trolox Antifade Reagent reduces phototoxicity and photobleaching of reagents to increase cell viability and prolong signal.

VectaCell Trolox Antifade Reagent

VectaCell Trolox Antifade Reagent is an antifading additive for live cell imaging. VectaCell Trolox Antifade Reagent contains both Trolox and its oxidized form Trolox-quinone. This redox system reduces photo-bleaching and blinking during live cell imaging.

Trolox is a water-soluble and cell-permeable analog of vitamin E that efficiently prevents formation of different reactive oxygen species, such as singlet oxygen (1O_2), superoxide anion (O_2^-) or hydrogen peroxide (H_2O_2). Photo-excitation of a fluorophore generates reactive oxygen species that can lead to photo-bleaching and oxidative damage in cells. Trolox has a cytoprotective effect and low cytotoxicity for different cell lines.



Product	Catalog Number
VectaCell™ Trolox Antifade Reagent	CB-1000

Accessory Reagents

VECTABOND Reagent Tissue Section Adhesive

VECTABOND Reagent chemically modifies the surface of glass to form a highly adherent charged surface. This charge significantly increases the adherence of both frozen and paraffin-embedded tissue sections and cell preparations to glass microscope slides and coverslips. Tissue sections will remain attached even when subjected to the most extreme conditions, such as high-temperature antigen retrieval and *in situ* hybridization. VECTABOND Reagent treated slides can be stored indefinitely.

ImmEdge Hydrophobic Barrier Pen

The ImmEdge Pen is a hydrophobic barrier (PAP) pen for immunohistochemistry and *in situ* hybridization. It provides a water-repellent barrier that keeps reagents localized on tissue specimens and prevents mixing of reagents when multiple sections are mounted on the same slide.

- Heat-stable
- Insoluble in alcohol and acetone
- Stable for use with buffers with and without detergent (Tween 20, Triton X-100, etc.)
- Completely removed by all commonly used xylene and xylene-substitute clearing agents
- Contains no ozone-depleting solvents
- Compatible with both enzyme- and fluorescence-based detection systems



ImmPrint Histology Pen

The ImmPrint Histology Pen is a permanent marking pen designed for writing on glass microscope slides, tissue cassettes, and most hard surfaces. Unlike other pens commonly used for histology, the ImmPrint Pen has a smooth writing tip that resists drying out.

- High-density, fast-drying, black ink
- Resistant to most organic solvents encountered in histological applications

Control Antibodies

These antibodies are IgG preparations for use as controls for primary antibodies made in rabbit, mouse, rat, or goat. Each has been purified from pooled serum of healthy adult animals and contain a spectrum of the IgG subclasses. When applied appropriately, these controls will help determine whether the primary antibody staining signal is specific for the antigen or whether staining is the result of non-specific adsorption of primary antibody to tissue sites.

Antigen Unmasking Solutions

Our Antigen Unmasking Solutions are highly effective at revealing antigens in formalin-fixed, paraffin-embedded tissue sections when used in combination with a high temperature treatment procedure. We offer two formulations of Antigen Unmasking Solution: Citrate-based solution (pH 6.0) and Tris-based solution (pH 9.0), each supplied as 100X concentrated stocks.

Product	Catalog Number
VECTABOND® Reagent (Tissue Section Adhesive)	SP-1800
ImmEdge® Hydrophobic Barrier PAP Pen	H-4000
ImmPrint™ Histology Pen	H-6100
Control Antibodies	
Rabbit IgG	I-1000
Mouse IgG	I-2000
Rat IgG	I-4000
Goat IgG	I-5000
Antigen Unmasking Solutions	
Citrate-based (100X) (pH 6.0)	H-3300
Tris-based (100X) (pH 9.0)	H-3301

Blocking Background Signal

Blocking agents minimize background signal from endogenous enzyme activity, biotin, and non-specific binding of tissue elements (charged particles, macromolecules, Fc receptors) with detection reagents. For IF applications special consideration should be given to the presence of autofluorescence.

TrueVIEW Autofluorescence Quenching Kits

TrueVIEW Autofluorescence Quenching Kits provide a novel way to remove unwanted fluorescence in tissue sections due to aldehyde fixation, red blood cells, and structural elements such as collagen and elastin. The unique formulation binds and effectively quenches the autofluorescent elements in even the most problematic tissues, such as kidney, spleen and pancreas.


The use of TrueVIEW Quenching reagent leads to significant enhancement in overall signal-to-noise in most immunofluorescence assays.

TrueVIEW Quenching reagent is a unique approach to diminish unwanted autofluorescence from non-lipofuscin sources, that retains the specific fluorescent antigen staining. The quenching action of the kit reagents therefore, provides the investigator with a clear, unambiguous, "true view" visualization of the intended target.

Why TrueVIEW Quencher?

- Specific reduction of autofluorescence from aldehyde fixation
- Improved signal-to-noise ratio
- Effective in even the most challenging tissues
- Easy-to-use, 5 minute incubation step
- Broad fluorophore compatibility
- Compatible with epifluorescence and confocal microscopes
- Antifade mounting medium included (with or without DAPI counterstain)



Serial sections of FFPE Human Prostate stained for epithelium (red)



*Both sections mounted with VECTASHIELD Vibrance Antifade Mounting Medium with DAPI.

"I would definitely use this reagent in the future - it is quick and reliable on multiple tissue types."

- Dr. K. Sadtler, Postdoctoral Fellow MIT Boston Children's Hospital



Adjacent human pancreas sections (FFPE) insulin using fluorescein label (green). Mounted with VECTASHIELD Vibrance Antifade Mounting Medium. Note significant reduction of autofluorescence in the TrueVIEW treated section (right) with the retention of specific staining.

BLOXALL® Endogenous Peroxidase and Alkaline Phosphatase Blocking Solution

BLOXALL is compatible with formalin-fixed, paraffin-embedded tissue sections, frozen tissue sections, and cell preparations. It is supplied ready-to-use in a convenient dropper bottle and only requires a brief 10-minute incubation.

Levamisole Solution

Specifically inhibits endogenous alkaline phosphatase activity that is added to the alkaline phosphatase substrate solution. It is supplied ready-to-use in a convenient dropper bottle.

Avidin/Biotin and Streptavidin/Biotin Blocking Kits

Both kits block all endogenous biotin and biotin receptors. Due to differing binding affinities and characteristics, kit selection is matched to the specific avidin or streptavidin detection system being used. Supplied ready-to-use in convenient dropper bottles.

Normal Sera and 2.5% Normal Sera

All our sera products are pooled samples collected from healthy adult animals, heat-treated and centrifuged to remove precipitates and then filtered. These sera are intended to be used for blocking non-specific binding or as an antibody diluent.

Bovine Serum Albumin (BSA)

Intended to be used as a diluent or a blocking agent and is free of impurities present in other grades of BSA which can introduce artifacts or increase background staining.

10x Casein Solution

A general blocking agent for IHC, nucleic acid blotting, protein blotting, and other applications.

Carbo-Free Blocking Solution

A protein-based agent that is essentially free of glycoproteins making it ideal for applications using lectins. Can be used to block non-specific binding or as an antibody diluent.

Animal-Free Blocker and Diluent, R.T.U.

A plant protein-derived solution intended for cell- and tissue-based IHC and IF applications. Can be used as an alternative to normal sera, BSA, casein and non-fat dry milk. Suitable for use with both HRP and AP enzyme conjugates and detection systems. Supplied as a ready-to-use solution, ideal in multiple antigen labeling IHC to streamline blocking.

Animal-Free Blocker (5x concentrate solution)

Similar to the R.T.U. format, this plant protein-derived blocking agent and diluent is an alternative to normal sera, BSA, casein and non-fat dry milk, however this concentrate is intended primarily for blotting applications.

Product	Catalog Number
Vector® TrueVIEW® Autofluorescence Quenching Kit	SP-8400
Vector® TrueVIEW® Autofluorescence Quenching Kit with DAPI	SP-8500
BLOXALL® Endogenous HRP/AP Blocking Solution	SP-6000
Levamisole Solution	SP-5000
Avidin/Biotin Blocking Kit	SP-2001
Streptavidin/Biotin Blocking Kit	SP-2002
Normal Goat Serum	S-1000
Normal Horse Serum	S-2000
Normal Chicken Serum	S-3000
Normal Swine Serum	S-4000
Normal Rabbit Serum	S-5000
2.5% Normal Goat Serum	S-1012
2.5% Normal Horse Serum	S-2012
Bovine Serum Albumin (BSA)	SP-5050
10x Casein Solution	SP-5020
Carbo-Free™ Blocking Solution	SP-5040
Animal-Free Blocker® and Diluent, R.T.U.	SP-5035
Animal-Free Blocker®	SP-5030

Contact Details

Ordering Information

Order online at vectorlabs.com

Orders may also be placed by email, telephone, or mail.

Please include the following with each order:

- Product name and catalog number
- Unit size and quantity
- Billing and shipping addresses
- Purchase order number
- Name, phone number, address and email address of person placing order

Orders using VISA, Mastercard, or American Express are accepted and processed immediately. Telephone orders over \$1000 may require written confirmation. A confirmation should be boldly marked "Confirming Order. Do Not Duplicate". Duplicate shipments due to incorrectly marked confirming orders cannot be returned for credit. No returned product will be accepted or credited without prior authorization from Vector Laboratories.

Please contact us to discuss discounts for custom or large orders.

Payment/shipping terms

For non-credit card orders, our payment terms are net 30 days from date of invoice, title and risk of loss transfer Ex Works (Incoterms 2010) Seller's location, freight prepaid and added unless shipped on Buyer's account (FedEx, UPS, DHL). Buyers are required to submit a credit application before credit terms are extended. Orders placed before 3 pm Pacific Time on Monday through Friday (excluding holidays) are usually processed the same day they are received. Unless requested otherwise, all products are shipped 2nd Day Air.



US Office

USA Headquarters:
Vector Laboratories, Inc.
6737 Mowry Avenue
Newark, CA 94560
Tel: +1 (650) 697-3600

Ordering and Technical Service (USA): +1 (800) 227-6666

Customer Service: vector@vectorlabs.com

Technical Support: technical@vectorlabs.com

International Inquiries: techintl@vectorlabs.com

Although provided in a highly purified form, our products are not intended for clinical diagnosis or drug use, nor have they been packaged under sterile conditions. All products listed in this catalog are for research purposes only. The listing of any product in this catalog does not imply the absence of a patent covering its use, does not constitute license under any existing or pending patent, nor is it intended or implied as a recommendation for the use of such products in infringement of any patent. The responsibility for determining the existence of such patents rests solely with the user.

Animal-Free Blocker, BLOXALL, Carbo-Free, HardSet, ImmEDGE, ImmPRESS, ImmPRINT, M.O.M., VECTABOND, VectaCell, VectaFluor, VECTASHIELD, VECTASHIELD VIBRANCE, and Vector are trademarks of Vector Laboratories, Inc
DyLight and Texas Red are trademarks of Pierce Biotechnology, Inc. All other trademarks cited herein are the property of their respective owners.

For more information please visit

vectorlabs.com


©2022 Vector Laboratories. All rights reserved.

For research use only. Not intended for animal or human therapeutic or diagnostic use.

LIT3002.Rev00



vectorlabs.com



—
Together we
breakthrough™



*Immunohistochemistry • Immunofluorescence
Lectins & Glycobiology • Bioconjugation*