




# Click Chemistry Toolbox<sup>2025</sup>


- ✱ Nascent Protein Synthesis Assay Kits
- ✱ Fluorescent Probes for Cu-free Click Chemistry
- ✱ Fluorogenic Azides Probes
- ✱ Azide Plus – Next Generation Azide Probes
- ✱ Iso-TaG Reagents and Kits
- ✱ Cleavable Click Chemistry Biotin Probes
- ✱ Click Chemistry Enrichment Kits and Media





● Click Chemistry Reaction Buffer Kits .....	5
Click Chemistry Auxiliary Reagents .....	6
● Click Chemistry Imaging Kits .....	7
● Nascent Protein Synthesis Assay Kits .....	8
● Fluorogenic Azide Probes .....	9
● Iso-TaG Reagents and Kits .....	10-11
● Azide Plus Reagents .....	12-15
● Fluorescent Azides .....	16-17
● Biotin Probes for Click Chemistry .....	18-19
● Fluorescent Alkyne Probes .....	20
● Fluorescent Probes for Copper-free Click Chemistry .....	22
Biotin Probes and Labeling Reagents for Copper-free Click Chemistry .....	23
● Enrichment Media and Kits, (Biotin-Streptavidin free) .....	24-27
● Enrichment Kits (Biotin-Streptavidin-based) .....	28-31
● Cleavable Click Chemistry Biotin Probes .....	30
DADPS Biotin Probes .....	30
Photocleavable Biotin Probes .....	31
Dde Biotin Probes .....	32
Diazo Biotin Probes .....	33
● Metabolic Labeling Reagents .....	34-37

# Copper-Catalyzed Click Reaction



**Figure 1** Schematic representation of copper-catalyzed click reaction.

Click reactions are defined more broadly as those that meet the necessary criteria of being selective, high yielding, wide in scope and having good reaction kinetics. These reactions possess extreme selectivity and biocompatibility, such that their participating reagents can form covalent bonds within richly functionalized biological systems—in some cases, living organisms.

Among many click reactions described up to date, the most widely used reaction is the Huisgen 1,3-dipolar cycloaddition of alkynes to azides to form 1,4-disubstituted-1,2,3-triazoles (**Figure 1**). The copper(I)-catalyzed azide-alkyne cycloaddition reaction (CuAAC) is mild and very efficient, requiring no protecting groups, and requiring no purification in many cases. The azide and alkyne functional groups are largely inert towards biological molecules and aqueous environments. Unlike other labels, the azide- and alkyne-tags are small enough that tagged biomolecules (e.g., azide- or alkyne-containing sugars, amino acids and nucleotides) are acceptable substrates for the enzymes that incorporate these building blocks into biopolymers such as proteins, DNA and RNA. This unique property paved the way for a very powerful, innovative and simple two-step labeling procedure. In the first step, an azide- or alkyne-containing biomolecule is actively incorporated into the protein. The second step, the detection step, uses the chemoselective ligation or “click” reaction between an azide and an alkyne. In the click reaction, the modified protein is detected with a corresponding azide- or alkyne-containing dye or hapten.

This powerful two-step procedure enables a large number of applications, such as detection of global RNA/DNA synthesis temporally and spatially in cells and tissues; detection and characterization of newly synthesized proteins; changes in spatial or temporal protein expression patterns; protein degradation resulting from disease, drug treatments, or environmental changes; visualization and characterization of various post-translational modifications (e.g. glycosylation, acylation, phosphorylation); and imaging bacterial cell wall biosynthesis.

### Click-&-Go® Click Chemistry Reaction Buffer Kit

The Click-&-Go® Click Chemistry Reaction Buffer Kit provides researchers — who have biomolecules labeled with an azide or alkyne and the corresponding click detection reagent — with all of the necessary reagents to perform a copper-catalyzed ligation reaction. Sufficient materials are provided to perform up to 25 copper-catalyzed click reactions for subsequent analysis by gel electrophoresis, western blot or mass spectrometry.



Product #	Description	Pkg. Size
CCT-1001	Click-&-Go® Click Chemistry Reaction Buffer Kit	1 kit

### Click-&-Go® Cell Reaction Buffer Kit

The Click-&-Go® Cell Reaction Buffer Kit provides researchers with everything required to perform a click reaction on cells tagged with an azide or alkyne with the corresponding click detection reagent for subsequent downstream analysis.

The performance and components of this kit are identical to Click-iT® Protein Reaction Buffer Kit from Thermo Fisher Scientific (Cat# C10269).

Product #	Description	Pkg. Size
CCT-1263	Click-&-Go® Cell Reaction Buffer Kit	1 kit


### Click-&-Go® Protein Reaction Buffer Kit

The Click-&-Go® Protein Reaction Buffer Kit provides researchers with everything required to perform a click reaction on azide or alkyne tagged proteins with the corresponding click detection reagent for subsequent downstream analysis.


The performance and components of this kit are identical to Click-iT® Protein Reaction Buffer Kit from Thermo Fisher Scientific (Cat# C10276).

Product #	Description	Pkg. Size
CCT-1262	Click-&-Go® Protein Reaction Buffer Kit	1 kit


THPTA		Catalog#	Unit
CAS:	760952-88-3	CCT-1010-100	100 mg
MW:	434.50	CCT-1010-500	500 mg
Solubility:	Water, DMSO, DMF	CCT-1010-1000	1000 mg
Description:	Water-soluble chelating agent	CCT-1010-5G	5 g




BTAA		Catalog#	Unit
CAS:	1334179-85-9	CCT-1236-100	100 mg
MW:	430.52	CCT-1236-500	500 mg
Solubility:	Water, DMSO, DMF	CCT-1236-1000	1000 mg
Description:	Water-soluble chelating agent		




BTES		Catalog#	Unit
CAS:	n/a	CCT-1237-100	100 mg
MW:	494.62	CCT-1237-500	500 mg
Solubility:	Water, DMSO, DMF	CCT-1237-1000	1000 mg
Description:	Water-soluble chelating agent		



BTTP		Catalog#	Unit
CAS:	n/a	CCT-1414-100	100 mg
MW:	430.56	CCT-1414-500	500 mg
Solubility:	DMSO, DMF, MeOH	CCT-1414-1000	1000 mg
Description:	Water-soluble chelating agent		



TBTA		Catalog#	Unit
CAS:	510758-28-8	CCT-1061-100	100 mg
MW:	530.62	CCT-1061-500	500 mg
Solubility:	Water, DMSO, DMF	CCT-1061-1000	1000 mg
Description:	Chelating agent		





Click-&Go® Plus Imaging Kits

The Click-&Go® Plus Imaging Kit is a general purpose imaging kit that is designed to perform a high sensitivity imaging of moderate-to-low abundance alkyne-containing biomolecules. The labeling kit utilizes the latest generation of copper-chelating azide capable of forming strong, active copper complexes that react almost instantaneously with alkynes under diluted conditions.

Each Click-&Go® Plus Imaging Kit includes the fluorescent azide plus probe and all of the reagents required to create a reaction cocktail.

Product #	Description	Pkg. Size
CCT-1313	Click-&Go® Plus 405 Imaging Kit	1 kit
CCT-1314	Click-&Go® Plus 488 Imaging Kit	1 kit
CCT-1315	Click-&Go® Plus 532 Imaging Kit	1 kit
CCT-1316	Click-&Go® Plus 546 Imaging Kit	1 kit
CCT-1317	Click-&Go® Plus 555 Imaging Kit	1 kit
CCT-1318	Click-&Go® Plus 568 Imaging Kit	1 kit
CCT-1319	Click-&Go® Plus 594 Imaging Kit	1 kit
CCT-1320	Click-&Go® Plus 647 Imaging Kit	1 kit






Click-&-Go® Plus OPP Protein Synthesis Assay Kits

Click-&-Go® Plus OPP Protein Synthesis Assay Kits enable fast, sensitive, and non-radioactive detection of protein synthesis using fluorescence microscopy or high-content imaging. O-propargyl-puromycin (OPP), an analog of puromycin that contains a terminal alkyne group, enters the acceptor site of ribosomes and incorporates into nascent polypeptide chains. OPP is not an amino acid analog, thus, OPP can be added directly to cells in complete media (i.e., methionine-containing) or used to detect in vivo protein synthesis. OPP that is incorporated into newly translated proteins is detected with fluorescent azides through a fast, highly-specific, and mild click reaction.

Product #	Description	Pkg. Size
CCT-1492	Click-&-Go® Plus 405 OPP Protein Synthesis Assay Kit	1 kit
CCT-1493	Click-&-Go® Plus 488 OPP Protein Synthesis Assay Kit	1 kit
CCT-1494	Click-&-Go® Plus 555 OPP Protein Synthesis Assay Kit	1 kit
CCT-1495	Click-&-Go® Plus 594 OPP Protein Synthesis Assay Kit	1 kit
CCT-1496	Click-&-Go® Plus 647 OPP Protein Synthesis Assay Kit	1 kit

O-propargyl-puromycin (OPP)

CAS: 1416561-90-4  
MW: 495.54  
Solubility: DMSO, DMF, water (pH adjusted to 5.0)




Catalog#	Unit
CCT-1407-5	5 mg
CCT-1407-25	25 mg
CCT-1407-100	100 mg



# CalFluor Azide Probes

A major shortcoming of the visualization of alkyne-tagged biomolecules with fluorescent azide probes through CuAAC reactions is the need to remove unreacted fluorescent probes. This is particularly problematic when imaging the intracellular environment, tissues of living organisms, or visualizing biomolecules in vivo. The difficulty of removing all unreacted fluorescent probes is also one of the major contributors to background signal and non-specific binding.


To overcome this shortcoming, the Carolyn Bertozzi group has designed fluorogenic azide probes that are activated by Cu-catalyzed or metal-free click chemistry. These azide probes are not fluorescent until they react with alkynes. Termed the CalFluors, these probes possess emission maxima that range from green to far-red wavelengths, and enable sensitive biomolecule detection under no-wash conditions. A number of reports showed that CalFluor probes are an indispensable tool for sensitive visualization of metabolically labeled molecules (glycans, DNA, RNA, and proteins) in cells, developing zebrafish, and mouse brain tissue slices under no-wash conditions.



# IsoTaG Reagents and Kits

A popular strategy for protein identification is the bottom-up shotgun proteomics approach. In this method, a mixture of proteins is subjected to proteolytic digestion, the resulting peptides are separated by LC and detected by MS, and their parent proteins are inferred from the assigned peptide sequences.

To convert MS data acquired from proteolytic digests into protein identifications, tandem MS can be used to obtain sequence information for individual peptides, followed by comparing an in-silico proteolytic digest of an organism's proteome. Typically, only the most abundant peptides are selected for fragmentation (**Figure 4**), whereas data for those peptides in relatively low quantities are not obtained. An inherent problem in shotgun proteomics is identifying proteins of low abundance, such as biomarkers for disease states, against a background of proteins whose concentrations can span up to 12 orders of magnitude.




**Figure 4** Traditional proteomics and Iso-TaG-directed proteomics workflow.

To address the unique challenges of identifying proteins of low abundance, a mass-independent chemical proteomics platform, termed *isotope targeted glycoproteomics* (IsoTaG), was developed by the Carolyn Bertozzi group. The platform is comprised of four central components: (i) metabolic labeling with a chemically functionalized glycan, (ii) chemical tagging and enrichment using an isotopic recoding affinity probe, (iii) directed tandem MS, and (iv) targeted glycopeptide assignment (**Figure 4**).

IsoTaG is performed by isotopic recoding and enrichment of metabolically labeled glycoproteins followed by directed tandem MS (MS2 or MSn) analysis and intact glycopeptide assignment. Isotopic recoding is accomplished by metabolic labeling of cell or tissue samples with azide- or alkyne-functionalized sugars, followed by chemical conjugation with a biotin probe bearing a unique isotopic signature.

In order to perform isotopic tagging, two IsoTaG probes encoded by zero [M] and two [M + 2] deuterium atoms are required. Probes with different encoding can be used and can be provided by Vector Laboratories though custom synthesis. The IsoTaG probe with zero, and that with two deuterium atoms [M, M + 2], can be used in different proportions; 1:1, 1:2, 1:3 and 1:4. Pattern recognition with isotopic ratio of 1:3 showed the highest fidelity.



**Figure 5** Cleavable IsoTaG probe encoded by zero deuterium atoms [M] (R = H) and two deuterium atoms [M+2] (R = D).

Through these probes, a unique isotopic signature is embedded exclusively into the (glyco)peptides. The isotopic signature serves as a computationally recognizable full-scan MS reporter. A computational algorithm, termed *isotopic signature transfer and mass pattern prediction* (IsoStamp), for the detection of recoded species in full-scan mass spectra, was also developed by the Carolyn Bertozzi group. IsoStamp compares observed and predicted isotopic envelopes to identify chemically tagged species in full-scan mass spectra.

IsoTaG has the potential to enhance any proteomics platform that employs chemical labeling for targeted protein identification, including isotope-coded affinity tagging, isobaric tagging for relative and absolute quantitation, and chemical tagging strategies for post-translational modification.

Product #	Description	Pkg. Size
CCT-1448	Click-&-Go® IsoTag Kit for Intact Glycopeptides Profiling *azide modified proteins*	1 kit
CCT-1449	Click-&-Go® IsoTag Kit for Intact Glycopeptides Profiling *alkyne modified proteins*	1 kit
CCT-1450	DADPS H2/D2 Biotin Azide, 2 mg each	1 set
CCT-1451	DADPS H2/D2 Biotin Alkyne, 2 mg each	1 set
CCT-1501	IsoTaG Biotin Azide Pack, 2 mg each	1 set
CCT-1502	IsoTaG Biotin Alkyne Pack, 2 mg each	1 set


#### Selected References:

- Woo, C.M., *et al.* (2015). Isotope-targeted glycoproteomics (IsoTaG): a mass-independent platform for intact N- and O-glycopeptide discovery and analysis. *Nat Methods*, **12**: 561–7.
- Woo, C. M., *et al.* (2017). Development of IsoTaG, a Chemical Glycoproteomics Technique for Profiling Intact N- and O-Glycopeptides from Whole Cell Proteomes. *J. Proteome Res.*, **16**: 1706–18.
- Gao, G., *et al.* (2017). Small Molecule Interactome Mapping by Photoaffinity Labeling Reveals Binding Site Hotspots for the NSAIDs. *J. Am. Chem. Soc.*, **140**: 4259–68.


Iso-Tag products are covered by U.S. Patent No.: 10,114,026. This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays, where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated, then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.

# Next Generation Azide Probes

Recent advances in the design of copper-chelating ligands, such as THPTA or BTAA that stabilize the Cu(I) oxidation state in aqueous solution, improve the kinetics of the copper-catalyzed azide-alkyne cycloaddition (CuAAC) reaction and greatly increase the sensitivity of alkyne detection. Copper-chelating ligands have also been shown to increase the biocompatibility of the CuAAC reaction by preventing the copper ions from causing biological damage<sup>1</sup>. The next step in improving the CuAAC reaction was the development of copper-chelating azides as more reactive substrates. Since it is speculated that the Cu(I)-azide association is the rate-determining step in the CuAAC catalytic cycle<sup>2</sup>, the introduction of a copper-chelating moiety at the azide reporter molecule allows for a dramatic raise of the effective Cu(I) concentration at the reaction site, enhancing the weakest link in the reaction rate acceleration (**Figure 7**). It has been proposed that the high reactivity of chelating azides comes from the rapid copper-azido group interaction which occurs prior to Cu(I) acetylide formation, and this renders the deprotonation of alkyne in the rate-determining step<sup>3</sup>. This concept was successfully exploited to perform CuAAC reactions using pyridine-based copper-chelating azides (picolyl azides) as substrates<sup>4-6</sup>. Nevertheless, the copper-chelating motif of picolyl azide molecules is not complete, requiring the presence of a copper chelator (e.g. THPTA) to achieve significant improvement in the kinetics of the CuAAC reaction<sup>3,4</sup>.




**Figure 6** Kinetic comparison of chelating azide and non-chelating conventional azide



**Figure 7**

In efforts to improve the performance of the CuAAC reaction in complex media, Vector Laboratories developed new chelating azides with a complete copper-chelating system in their structure, termed “Azides Plus” (**Figure 8**). These azides are capable of forming strong, active copper complexes and are therefore considered both reactant and catalyst in the CuAAC reaction. Using these types of azides, the CuAAC reaction becomes a bimolecular reaction and displays much faster kinetics compared to the CuAAC reaction performed with conventional azides.



**Figure 8**






Figure 9


Comparative kinetic measurements for the CuAAC reaction (**Figure 9**) were performed using an agarose-alkyne resin labeling experiment (3.0 mM CuSO<sub>4</sub>, with (6.0 mM) or without THPTA ligand) using Cy5 Azide Plus, Cy5 Picolyl Azide, and Cy5 bis-Triazole Azide – the fastest copper-chelating azide that has been reported to date<sup>7</sup>. As expected, the picolyl azide containing the incomplete copper-chelating motif displays relatively slow reactivity, in particular without the presence of THPTA. The kinetic data shows that completing a copper-chelating moiety greatly enhances reactivity, and importantly does not require the presence of copper-chelating ligands. Interestingly, the copper-chelating azides developed by Vector Laboratories display almost identical reactivity in the CuAAC reaction compared to the most reactive copper-chelating azide reported up to now<sup>7</sup>, bis-triazole azide.



Cy5 Azide Plus



Cy5 bis-Triazole Azide



Cy5 Picolyl Azide

The new copper chelating azides allow the formation of azide copper complexes that react almost instantaneously with alkynes under diluted conditions. This unprecedented reactivity in the CuAAC reaction is of special value for the detection of low abundance targets, improving biocompatibility, and any other application where greatly improved S/N ratio is highly desired.

#### Selected References:

- Hong, V., *et al.* (2010). Labeling Live Cells by Copper-Catalyzed Alkyne – Azide Click Chemistry. *Bioconjugate Chem.*, **21**, 1912–6.
- Rodionov, V.O., *et al.* (2007). Ligand-accelerated Cu-catalyzed azide-alkyne cycloaddition: A mechanistic report. *J. Am. Chem. Soc.*, **129**, 12705–12. Presolski, S.I., *et al.* (2010). Tailored ligand acceleration of the cu-catalyzed azide-alkyne cycloaddition reaction: Practical and mechanistic implications. *J. Am. Chem. Soc.*, **132**, 14570–6.
- Kuang, G.-C., *et al.* (2011). Experimental investigation on the mechanism of chelation-assisted, copper (ii) acetate-accelerated azide-alkyne cycloaddition. *J. Am. Chem. Soc.*, **133**, 13984–4001.
- Jiang, H., *et al.* (2014). Monitoring Dynamic Glycosylation in Vivo Using Supersensitive Click Chemistry. *Bioconjugate Chem.*, **25**: 698-706.
- Uttamapinant, C., *et al.* (2012). Fast, Cell-Compatible Click Chemistry with Copper-Chelating Azides for Biomolecular Labeling. *Angew. Chem. Int. Ed.*, **51**: 5852-6.
- Gaebler A., *et al.* (2016). A highly sensitive protocol for microscopy of alkyne lipids and fluorescently tagged or immunostained proteins. *J. Lipid. Res.*, **57**:1934-47.
- Bevilacqua, V., *et al.* (2014). Copper-Chelating Azides for Efficient Click Conjugation Reactions in Complex Media. *Angew. Chem. Int. Ed.*, **53**, 5872-6.

## Fluorescent Azide Plus Probes

Vector Laboratories offers a wide section of fluorescent Azide Plus probes, including AZDyes, Cy Dyes and classic dyes conjugated to azide groups. The photophysical properties of our AZDyes are an exact match to Alexa Fluor® Dyes. The combination of the exceptional reactivity of the azide plus moiety, biocompatibility and brightness of the AZDyes makes these probes of special value not only for the detection of low abundance targets, but also for all other applications where increased S/N ratio is of great value.


Description	Ex/Em	Emission Color	Pkg. Size	Product #
AZDye 350 Azide Plus	346/445	Blue	1 mg	CCT-1477-1
			5 mg	CCT-1477-5
			25 mg	CCT-1477-25
AZDye 405 Azide Plus	402/424	Blue	1 mg	CCT-1474-1
			5 mg	CCT-1474-5
			25 mg	CCT-1474-25
AZDye 488 Azide Plus	494/517	Green	1 mg	CCT-1475-1
			5 mg	CCT-1475-5
			25 mg	CCT-1475-25
AZDye 532 Azide Plus	532/554	Orange	1 mg	CCT-1476-1
			5 mg	CCT-1475-5
			25 mg	CCT-1475-25
AZDye 546 Azide Plus	554/570	Orange	1 mg	CCT-1478-1
			5 mg	CCT-1478-5
			25 mg	CCT-1478-25
AZDye 555 Azide Plus	555/572	Red	1 mg	CCT-1479-1
			5 mg	CCT-1479-5
			25 mg	CCT-1479-25
AZDye 568 Azide Plus	578/602	Red	1 mg	CCT-1480-1
			5 mg	CCT-1480-5
			25 mg	CCT-1480-25
AZDye 594 Azide Plus	590/617	Red	1 mg	CCT-1481-1
			5 mg	CCT-1481-5
			25 mg	CCT-1481-25
AZDye 647 Azide Plus	648/671	Near IR	1 mg	CCT-1482-1
			5 mg	CCT-1482-5
			25 mg	CCT-1482-25

Alexa Fluor® is a registered trademark of Thermo Fisher Scientific.

Description	Ex/Em	Emission Color	Pkg. Size	Product #
PB Azide Plus (Pacific Blue®equivalent)	410/455	Blue	1 mg	CCT-1483-1
			5 mg	CCT-1483-5
			25 mg	CCT-1483-25
TAMRA Azide Plus	553/575	Orange	1 mg	CCT-1486-1
			5 mg	CCT-1486-5
			25 mg	CCT-1486-25
Cy3 Azide Plus	555/572	Red	1 mg	CCT-1484-1
			5 mg	CCT-1484-5
			25 mg	CCT-1484-25
Cy5 Azide Plus	647/663	Near IR	1 mg	CCT-1485-1
			5 mg	CCT-1485-5
			25 mg	CCT-1485-25

### Biotin Azide Plus


**CAS:** n/a  
**MW:** 582.72  
**Solubility:** DMSO, DMF  
**Description:** Biotinylation reagent with superior kinetics in copper-catalyzed click reactions.



Catalog#	Unit
CCT-1488-1	1 mg
CCT-1488-5	5 mg
CCT-1488-25	25 mg
CCT-1488-100	100 mg

### Dde Biotin Azide Plus

**CAS:** n/a  
**MW:** 815.98  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Biotinylation reagent with superior kinetics in copper-catalyzed click reactions.



Catalog#	Unit
CCT-1489-1	1 mg
CCT-1489-5	5 mg
CCT-1489-25	25 mg



## Fluorescent Azides

Vector Laboratories offers the largest selection of fluorescent azide probes for click chemistry. Our selection of fluorescent probes includes AZDyes, Cy Dyes and classic dyes conjugated to azide groups. The photophysical properties of our AZDyes are an exact match to Alexa Fluor® Dyes.

Description	Ex/Em	Emission Color	Pkg. Size	Product #
AZDye 350 Azide	346/445	Blue	1 mg	CCT-1267-1
			5 mg	CCT-1267-5
			25 mg	CCT-1267-25
AZDye 405 Azide	402/424	Blue	1 mg	CCT-1307-1
			5 mg	CCT-1307-5
			25 mg	CCT-1307-25
AZDye 488 Azide	494/517	Green	1 mg	CCT-1275-1
			5 mg	CCT-1275-5
			25 mg	CCT-1275-25
AZDye 532 Azide	532/554	Orange	1 mg	CCT-1279-1
			5 mg	CCT-1279-5
			25 mg	CCT-1279-25
AZDye 546 Azide	554/570	Orange	1 mg	CCT-1283-1
			5 mg	CCT-1283-5
			25 mg	CCT-1283-25
AZDye 555 Azide	555/572	Red	1 mg	CCT-1287-1
			5 mg	CCT-1287-5
			25 mg	CCT-1287-25
AZDye 568 Azide	578/602	Red	1 mg	CCT-1291-1
			5 mg	CCT-1291-5
			25 mg	CCT-1291-25
AZDye 594 Azide	590/617	Red	1 mg	CCT-1295-1
			5 mg	CCT-1295-5
			25 mg	CCT-1295-25
AZDye 647 Azide	648/671	Near IR	1 mg	CCT-1299-1
			5 mg	CCT-1299-5
			25 mg	CCT-1299-25

Alexa Fluor® is a registered trademark of Thermo Fisher Scientific.


Visit [vectorlabs.com](https://vectorlabs.com) for more detailed information

Description	Ex/Em	Emission Color	Pkg. Size	Product #
<b>OG 488 Azide</b> Replacement of Invitrogen's Oregon Green® 488 Azide (Catalog number: O10180)	496/524	Green	1 mg	CCT-1264-1
			5 mg	CCT-1264-5
			25 mg	CCT-1264-25
			100 mg	CCT-1264-100
<b>Carboxyrhodamine 110 Azide</b>	501/523	Green	1 mg	CCT-AZ105-1
			5 mg	CCT-AZ105-5
			25 mg	CCT-AZ105-25
			100 mg	CCT-AZ105-100
<b>TAMRA Azide</b>	553/575	Orange	1 mg	CCT-AZ109-1
			5 mg	CCT-AZ109-5
			25 mg	CCT-AZ109-25
			100 mg	CCT-AZ109-100
<b>5-TAMRA Azide</b> Replacement of Invitrogen's Tetramethylrhodamine Azide (Catalog number: T10182).	553/575	Orange	1 mg	CCT-1245-1
			5 mg	CCT-1245-5
			25 mg	CCT-1245-25
			100 mg	CCT-1245-100
<b>Cy3 Azide</b>	553/569	Red	1 mg	CCT-AZ119-1
			5 mg	CCT-AZ119-5
			25 mg	CCT-AZ119-25
			100 mg	CCT-AZ119-100
<b>Cy5 Azide</b>	649/671	Near IR	1 mg	CCT-AZ118-1
			5 mg	CCT-AZ118-5
			25 mg	CCT-AZ118-25
			100 mg	CCT-AZ118-100
<b>Cy5.5 Azide</b>	678/694	Near IR	1 mg	CCT-1059-1
			5 mg	CCT-1059-5
			25 mg	CCT-1059-25
			100 mg	CCT-1059-100
<b>Cy7 Azide</b>	753/775	Near IR	1 mg	CCT-1052-1
			5 mg	CCT-1052-5
			25 mg	CCT-1052-25
			100 mg	CCT-1052-100

## Biotin Azide

CAS: n/a  
MW: 615.79  
Solubility: DMSO, DMF, MeOH  
Description:

Exact replacement of  
Invitrogen's Biotin Azide  
(PEG4 carboxamide-6-Azido-hexanyl Biotin),  
Catalog number: B10184




Catalog#

Unit

CCT-1265-5 5 mg  
CCT-1265-25 25 mg  
CCT-1265-100 100 mg

## Biotin-PEG3-Azide

CAS: 875770-34-6  
MW: 444.55  
Solubility: DMSO, DMF  
Description: Biotinylation reagent




Catalog#

Unit

CCT-AZ104-5 5 mg  
CCT-AZ104-25 25 mg  
CCT-AZ104-100 100 mg  
CCT-AZ104-1000 1000 mg

## Biotin Azide Plus

CAS: n/a  
MW: 582.72  
Solubility: DMSO, DMF, MeOH  
Description: Biotinylation reagent with superior kinetics in copper-catalyzed click reactions.




Catalog#

Unit

CCT-1488-1 1 mg  
CCT-1488-5 5 mg  
CCT-1488-25 25 mg  
CCT-1488-100 100 mg

## Biotin Picolyl Azide

CAS: n/a  
MW: 622.74  
Solubility: DMSO, DMF  
Description: Biotinylation reagent with superior kinetics in copper-catalyzed click reactions.




Catalog#

Unit

CCT-1167-5 5 mg  
CCT-1167-25 25 mg  
CCT-1167-100 100 mg

## Dde Biotin Picolyl Azide

CAS: n/a  
MW: 815.98  
Solubility: DMSO, DMF, THF, DCM  
Description: Biotinylation reagent with superior kinetics in copper-catalyzed click reactions.



Catalog#


Unit

CCT-1186-1 1 mg  
CCT-1186-5 5 mg  
CCT-1186-25 25 mg

### Biotin Alkyne

CAS: n/a  
MW: 528.26  
Solubility: DMSO, DMF


Description: [Exact replacement of Invitrogen's Biotin Alkyne \(PEG4 carboxamide-Propargyl Biotin\), Catalog number: B10185](#)



Catalog#	Unit
CCT-1266-5	5 mg
CCT-1266-25	25 mg
CCT-1266-100	100 mg

### Biotin-PEG4-Alkyne

CAS: 1262681-31-1  
MW: 457.58  
Solubility: DMSO, DMF, MeOH  
Description: Biotinylation reagent




Catalog#	Unit
CCT-TA105-5	5 mg
CCT-TA105-25	25 mg
CCT-TA105-100	100 mg
CCT-TA105-1000	1000 mg

## Biotin Probes for Cu-free Click Chemistry

### DBCO-PEG4-Biotin


CAS: 1255942-07-4  
MW: 749.92  
Solubility: DMSO, DMF, THF, MeOH  
Description: Biotinylation reagent



Catalog#	Unit
CCT-A105-5	5 mg
CCT-A105-25	25 mg
CCT-A105-100	100 mg

### WS DBCO-Biotin

CAS: 1363444-70-5  
MW: 653.77  
Solubility: DMSO, DMF, THF, DCM  
Description: Water-soluble biotinylation reagent



Catalog#	Unit
CCT-A116-1	1 mg
CCT-A116-5	5 mg
CCT-A116-25	25 mg
CCT-A116-100	100 mg

## Fluorescent Alkyne Probes


Vector Laboratories offers a wide section of fluorescent alkyne probes covering the entire UV-Vis spectrum. Our selection of fluorescent probes includes AZDyes, Cy Dyes and classic dyes conjugated to terminal alkynes. The photophysical properties of our AZDyes are an exact match to Alexa Fluor® Dyes.

Description	Ex/Em	Emission Color	Pkg. Size	Product #
AZDye 350 Alkyne	346/445	Blue	1 mg	CCT-1269-1
			5 mg	CCT-1269-5
			25 mg	CCT-1269-25
AZDye 405 Alkyne	402/424	Blue	1 mg	CCT-1309-1
			5 mg	CCT-1309-5
			25 mg	CCT-1309-25
AZDye 488 Alkyne	494/517	Green	1 mg	CCT-1277-1
			5 mg	CCT-1277-5
			25 mg	CCT-1277-25
AZDye 532 Alkyne	532/554	Orange	1 mg	CCT-1281-1
			5 mg	CCT-1281-5
			25 mg	CCT-1281-25
AZDye 546 Alkyne	554/570	Orange	1 mg	CCT-1285-1
			5 mg	CCT-1285-5
			25 mg	CCT-1285-25
AZDye 555 Alkyne	554/569	Red	1 mg	CCT-1289-1
			5 mg	CCT-1289-5
			25 mg	CCT-1289-25
AZDye 568 Alkyne	578/602	Red	1 mg	CCT-1293-1
			5 mg	CCT-1293-5
			25 mg	CCT-1293-25
AZDye 594 Alkyne	590/617	Red	1 mg	CCT-1297-1
			5 mg	CCT-1297-5
			25 mg	CCT-1297-25
AZDye 647 Alkyne	648/671	Near IR	1 mg	CCT-1301-1
			5 mg	CCT-1301-5
			25 mg	CCT-1301-25

Visit [vectorlabs.com](https://vectorlabs.com) for a full list of fluorescent terminal alkynes

Alexa Fluor® is a registered trademark of Thermo Fisher Scientific.

# Cu-Free Click Chemistry




**Figure 10** Schematic representation of a SPAAC ligation reaction.

The strain-promoted alkyne-azide cycloaddition reaction, also termed the Cu-free click reaction, is a bioorthogonal reaction utilizing a pair of reagents – cyclooctynes and azides – that exclusively and efficiently react with each other while remaining inert to naturally occurring functional groups such as amines (**Figure 10**). SPAAC enables labeling of a wide variety of biomolecules without any auxiliary reagents in an aqueous and otherwise complex chemical environment through the formation of a stable triazole.

Among the large number of known cyclooctynes, the so-called DBCO (dibenzocyclooctynes) compounds comprise a class of reagents that possesses reasonably fast kinetics and good stability in aqueous buffers. Within physiological temperature and pH ranges, the DBCO group will not react with amines or hydroxyls that are naturally present in many biomolecules. Additionally, reaction of the DBCO group with the azide group is significantly faster than with sulfhydryl groups ( $-SH$ , thiol).

Unlike many other cyclooctynes, DBCO reagents possess an embedded chromophore that allows for the simple and non-destructive spectroscopic identification of DBCO-containing compounds. This chromophore can also be used for spectroscopic estimation of total incorporated DBCO molecules into a biopolymer.

Another important feature of DBCO compounds is that the progress of SPAAC ligation can be followed in real time by simple UV-Vis spectroscopy. As the “click reaction” progresses the signature absorbance band at 310 nm disappears as illustrated **Figure 11**.



**Figure 11** Progress of SPAAC ligation reaction followed by UV-Vis spectroscopy.

## Fluorescent Probes for Copper-free Click Chemistry

In applications where the presence of copper is a concern, probes that react with azides via a copper-free click chemistry reaction to form stable triazoles are an ideal alternative to copper-requiring fluorescent alkynes. We offer the largest selection of fluorescent probes for copper-less azide imaging, covering the entire UV-Vis spectrum. Our selection of fluorescent probes includes AZDyes, Cy Dyes and classic dyes conjugated to DBCO alkynes.


Description	Ex/Em	Emission Color	Pkg. Size	Product #
AZDye 405 DBCO	402/424	Blue	1 mg	CCT-1310-1
			5 mg	CCT-1310-5
			25 mg	CCT-1310-25
AZDye 430 DBCO	430/537	Green	1 mg	CCT-1274-1
			5 mg	CCT-1274-5
			25 mg	CCT-1274-25
AZDye 488 DBCO	494/517	Green	1 mg	CCT-1278-1
			5 mg	CCT-1278-5
			25 mg	CCT-1278-25
AZDye 532 DBCO	532/554	Orange	1 mg	CCT-1282-1
			5 mg	CCT-1282-5
			25 mg	CCT-1282-25
AZDye 546 DBCO	554/570	Orange	1 mg	CCT-1286-1
			5 mg	CCT-1286-5
			25 mg	CCT-1286-25
AZDye 555 DBCO	555/572	Red	1 mg	CCT-1290-1
			5 mg	CCT-1290-5
			25 mg	CCT-1290-25
AZDye 568 DBCO	578/602	Red	1 mg	CCT-1294-1
			5 mg	CCT-1294-5
			25 mg	CCT-1294-25
AZDye 594 DBCO	590/617	Red	1 mg	CCT-1298-1
			5 mg	CCT-1298-5
			25 mg	CCT-1298-25
AZDye 647 DBCO	648/671	Near IR	1 mg	CCT-1302-1
			5 mg	CCT-1302-5
			25 mg	CCT-1302-25

Visit [vectorlabs.com](https://vectorlabs.com) to browse a full list of fluorescent probes for Cu-free click chemistry.



## DBCO-PEG4-Biotin


**CAS:** 1255942-07-4  
**MW:** 749.92  
**Solubility:** DMSO, DMF, THF, MeOH  
**Description:** Biotinylation reagent



Catalog#	Unit
CCT-A105-5	5 mg
CCT-A105-25	25 mg
CCT-A105-100	100 mg

## WS DBCO-Biotin


**CAS:** 1363444-70-5  
**MW:** 653.77  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Water-soluble biotinylation reagent



Catalog#	Unit
CCT-A116-1	1 mg
CCT-A116-5	5 mg
CCT-A116-25	25 mg
CCT-A116-100	100 mg

## DBCO-Sulfo-NHS Ester


**CAS:** 1400191-52-7  
**MW:** 532.50  
**Solubility:** Water, DMSO, DMF  
**Description:** Water-soluble labeling reagent



Catalog#	Unit
CCT-A124-10	10 mg
CCT-A124-25	25 mg
CCT-A124-100	100 mg
CCT-A124-500	500 mg

## DBCO-PEG5-NHS Ester


**CAS:** 1378531-80-6  
**MW:** 693.74  
**Solubility:** DMSO, DMF, DCM, THF  
**Description:** Labeling reagent with improved stability and solubility



Catalog#	Unit
CCT-A102P-2	4x2 mg
CCT-A102P-10	10 mg
CCT-A102P-25	25 mg
CCT-A102P-100	100 mg
CCT-A102P-500	500 mg
CCT-A102P-1G	1 g

## Sulfo DBCO-Maleimide


**CAS:** n/a  
**MW:** 578.59  
**Solubility:** Water, DMSO, DMF, DCM, THF  
**Description:** Water-soluble, sulfhydryl-reactive labeling reagent



Catalog#	Unit
CCT-1230-10	10 mg
CCT-1230-25	25 mg
CCT-1230-100	100 mg
CCT-1230-500	500 mg

## Sulfo DBCO-PEG4-Maleimide

**CAS:** n/a  
**MW:** 825.89  
**Solubility:** Water, DMSO, DMF, DCM, THF  
**Description:** Water-soluble, sulfhydryl-reactive labeling reagent with hydrophilic PEG<sub>4</sub> spacer arm



Catalog#	Unit
CCT-1231-10	10 mg
CCT-1231-25	25 mg
CCT-1231-100	100 mg
CCT-1231-500	500 mg

Full list of DBCO reagents available at [vectorlabs.com](https://vectorlabs.com)

# Enrichment Media and Kits

## Click-&-Go® Protein Enrichment Kits (Biotin-Streptavidin Free)


The ability to detect and characterize newly synthesized proteins, changes in spatial or temporal protein expression patterns, or protein degradation resulting from disease, drug treatments, or environmental changes, is an important parameter in cytotoxicity measurements. In most published studies<sup>1</sup>, azide- or alkyne-metabolically labeled, newly synthesized proteins were isolated from a pre-existing pool of proteins by an in-solution click reaction with biotin-alkyne or biotin-azide followed by capture on streptavidin resin. It was reported that with using such a strategy, newly synthesized proteins comprised only 10-20% of the isolated proteins<sup>2</sup>. To address this shortcoming of biotin-streptavidin based enrichment Vector Laboratories has developed an enrichment protocol that allows for direct, covalent capture of alkyne/azide tagged proteins onto azide- or alkyne-modified agarose resin followed by stringent washes to remove nonspecific resin-bound proteins prior to digestion and LCMS analysis.

Another recent study<sup>3</sup> assessed the level of the non-labeled proteins in BONCAT samples by performing the alkyne-based BONCAT sample preparation using HEK-TrKB cells that were labeled with AHA and comparing these results to a control experiment in which the same sample preparation was performed with the same amount of lysate from unlabeled cells. Both the BONCAT and the control samples were analyzed by LCMS. This study consistently identified dramatically more peptides from the BONCAT samples (2371, 2578, and 2681) than from the control samples (69, 19 and 83) at 1% FDR. Moreover, the peptides from the control samples generally had very low signals compared to those from the BONCAT samples. This result shows that the alkyne resin-based enrichment method has minimal contamination from non-AHA-labeled proteins and can be used to isolate high-purity nascent proteomes.

Direct, covalent capture of azide- or alkyne-tagged proteins onto agarose resin represents a substantial improvement compared to the biotin tag-based approach. This is ideal for the covalent capture of specific sub-classes of proteins which have been metabolically, enzymatically, or chemically azido-or alkyne-tagged onto a resin via Cu(I)-catalyzed azide-alkyne cycloaddition (CuAAC). The resin containing the covalently attached proteins can be washed with high stringency, virtually eliminating any non-specifically bound proteins *without causing loss of target proteins*. Upon protease digestion, it yields a highly specific peptide pool that is ideal for mass spectroscopy (e.g., LC MS/MS) based analysis.

### References:

1. (a) Shen, W., *et al.* (2014) Acute synthesis of CPEB is required for plasticity of visual avoidance behavior in *Xenopus*. *Cell Rep.*, **6**: 737 – 47. (b) Lu, Y., *et al.* (2014) Prometastatic GPCR CD97 is a direct target of tumor suppressor microRNA-126. *ACS Chem. Biol.*, **9**: 334 – 8. (c) Eichelbaum, K., *et al.* (2012) Selective enrichment of newly synthesized proteins for quantitative secretome analysis. *Nat. Biotechnol.*, **30**: 984 – 90. (d) Bagert, J. D., *et al.* (2014) Quantitative, time-resolved proteomic analysis by combining bioorthogonal noncanonical amino acid tagging and pulsed stable isotope labeling by amino acids in cell culture. *Mol. Cell Proteomics*, **13**: 1352 – 8. (e) Choi, K. Y., *et al.* (2012) Defining TNF-alpha and IL-1beta induced nascent proteins: combining bioorthogonal non-canonical amino acid tagging and proteomics. *J. Immunol. Methods*, **382**: 189 – 95. (f) Hodas, J. J., *et al.* (2012) Dopaminergic modulation of the hippocampal neuropil proteome identified by bioorthogonal noncanonical amino acid tagging (BONCAT). *Proteomics*, **12**: 2464 – 76.
2. Howden, A. J., *et al.* (2013) QuaNCAT: quantitating proteome dynamics in primary cells. *Nat. Methods*, **10**: 343 – 6.
3. Zhang G., *et al.* (2014) In-Depth Quantitative Proteomic Analysis of de Novo Protein Synthesis Induced by Brain-Derived Neurotrophic Factor. *J. Proteome Res.*, **13**, 5707 – 14.



**Figure 12** Schematic representation of pull-down workflows for biotin-streptavidin based and biotin-streptavidin free enrichment protocols.



### Click-&-Go® Protein Enrichment Kits, non-cleavable

The Click-&-Go® Protein Enrichment Kit is an efficient tool for covalent capture of azido- or alkyne-tagged proteins on a alkyne- or azide-agarose resin. The kit contains specially formulated components to both catalyze the click reaction and prevent non-specific binding to the alkyne- or azide modified resins. The alkyne- or azide-modified proteins, or their post-translationally modified forms, are captured from complex protein extracts on the azide-alkyne resin supplied. Once covalently attached to the resin via copper-catalyzed click chemistry, the beads can be washed with the highest stringency, virtually eliminating any non-specifically bound proteins to yield a highly enriched population of nascent molecules. Upon protease digestion, this yields a highly pure peptide pool that is ideal for mass spectrometry (e.g., LC MS/MS) based analysis.

Product #	Description	Pkg. Size
CCT-1039	Click-&-Go Protein Enrichment Kit *for capture of alkyne-modified proteins*	1 kit
CCT-1033	Click-&-Go Protein Enrichment Kit *for capture of azide-modified proteins*	1 kit
CCT-1235	Click-&-Go Plus Protein Enrichment Kit *for capture of alkyne-modified proteins*	1 kit


## Click Functionalized Agarose

Product #	Description	Pkg. Size
Alkyne agarose resin, 50% slurry	2 mL	CCT-1032-2
	25 mL	CCT-1032-25
Azide agarose resin, 50% slurry	2 mL	CCT-1038-2
	25 mL	CCT-1038-25

Click Functionalized Magnetic Beads are also available, please visit [Vector Laboratories website](#)

## Click-&-Go® Dde Protein Enrichment Kits

The Dde linker is stable towards acidic or basic conditions, generally applied buffer systems, and reactive species that are present in a cell extract. It also can withstand harsh wash conditions in order to virtually eliminate any non-specifically bound proteins. The captured proteins can be chemoselectively released under mild aqueous buffered conditions with 2% hydrazine to yield a highly enriched population of intact proteins.



**Figure 14** Schematic representation of hydrazine induced release.

Product #	Description	Pkg. Size
CCT-1153	Click-&-Go Dde Protein Enrichment Kit *for capture and release of alkyne-modified proteins*	1 kit
CCT-1152	Click-&-Go Dde Protein Enrichment Kit *for capture and release of azide-modified proteins*	1 kit



Click-&-Go® Enrichment Kits for Click Chemistry

The Click-&-Go® Protein Capture Kit provides all of the necessary reagents to perform a conventional capture of azide- or alkyne modified proteins though click labeling with a biotin reagent followed by capture on high-capacity streptavidin agarose resin. The kit includes specially formulated components to perform copper-catalyzed click reactions and subsequent capture on high-capacity streptavidin agarose. Sufficient material is supplied for 25 enrichments based on the provided protocol. The kit provides azide/alkyne labeled BSA as a positive control.

Product #	Description	Pkg. Size
CCT-1441	Click-&-Go Protein Enrichment Kit *for capture of alkyne-modified proteins*	1 kit
CCT-1440	Click-&-Go Protein Enrichment Kit *for capture of azide-modified proteins*	1 kit

### Click-&-Go® DADPS Protein Enrichment Kit for Click Chemistry (acid cleavable)

The Click-&-Go® DADPS Protein Enrichment Kit provides all of the necessary reagents to perform enrichment of azide-modified proteins through conventional biotin-streptavidin affinity purification. The kit includes an acid cleavable DADPS Biotin linker that allows for the release of captured proteins for intact protein analysis or on-beads digestion followed by the release of peptides for subsequent downstream analysis by mass spectrometry. Captured biomolecules can be released under mild conditions, such as 5% aqueous formic acid. Sufficient materials are supplied for 25 enrichments based on the provided protocol below. The kit provides azide labeled BSA as a positive control.

Product #	Description	Pkg. Size
CCT-1443	Click-&-Go DADPS Protein Enrichment Kit *for capture of alkyne-modified proteins*	1 kit
CCT-1442	Click-&-Go DADPS Protein Enrichment Kit *for capture of azide-modified proteins*	1 kit


### Click-&-Go® Dde Protein Enrichment Kit for Click Chemistry (hydrazine cleavable)

The Click-&-Go® Dde Protein Enrichment Kit provides all of the necessary reagents to perform enrichment of alkyne-modified proteins through conventional biotin-streptavidin affinity purification. The kit includes a cleavable Dde Biotin linker that allows for the release of captured proteins for intact protein analysis or on-beads digestion followed by the release of peptides for subsequent downstream analysis by mass spectrometry. Captured biomolecules can be released under mild conditions, such as 2% aqueous hydrazine. Sufficient materials are supplied for 25 enrichments based on the provided protocol. The kit provides alkyne labeled BSA as a positive control.

Product #	Description	Pkg. Size
CCT-1445	Click-&-Go Dde Protein Enrichment Kit *for enrichment of alkyne-modified proteins*	1 kit
CCT-1444	Click-&-Go Protein Enrichment Kit *for enrichment of azide-modified proteins*	1 kit



## DADPS Biotin Probes




The extraordinary strength of the biotin-streptavidin interaction allows for efficient capturing of even highly dilute targets; however, it makes recovery of proteins from affinity resins challenging. Conventional methods to elute biotinylated proteins from immobilized avidin include the following: (i) denaturation of streptavidin by boiling the resin in a denaturing buffer that may include high concentrations of chaotropic salts, (ii) trypsin digestion of proteins while they are bound to the resin, or (iii) elution of proteins with excess free biotin. These protocols can co-elute contaminant proteins by releasing nonspecifically bound proteins and/or naturally biotinylated proteins concurrently with labeled proteins. In addition, some of these methods can cause elution of high levels of resin-based peptides along with the proteins of interest, resulting in further sample contamination.

DADPS (dialkoxydiphenylsilane) Biotin probes eliminate a major limitation of the biotin-streptavidin affinity purification. These reagents contain a biotin moiety linked to an azide moiety through a spacer arm containing a cleavable DADPS linker. Captured biomolecules can be efficiently released under mild conditions (5% or 10% formic acid, 0.5 h) and the small molecular fragment is left on the labeled protein following cleavage. These features make the DADPS probe especially attractive for use in biomolecular labeling and proteomic studies.

### DADPS Biotin Azide

**CAS:** n/a  
**MW:** 886.19  
**Solubility:** DMSO, DMF  
**Description:** Cleavable biotinylation reagent




**Catalog#** **Unit**

CCT-1330-1 1 mg  
 CCT-1330-5 5 mg  
 CCT-1330-25 25 mg

### DADPS Biotin Alkyne

**CAS:** n/a  
**MW:** 827.12  
**Solubility:** DMSO, DMF  
**Description:** Cleavable biotinylation reagent




**Catalog#** **Unit**

CCT-1331-1 1 mg  
 CCT-1331-5 5 mg  
 CCT-1331-25 25 mg

### Selected References:

1. Szychowski, J., et al. (2010). Cleavable Biotin Probes for Labeling of Biomolecules via Azide-Alkyne Cycloaddition. *J. Am. Chem. Soc.*, **132**: 18351.
2. Jinxu, G., et al. (2012). Small Molecule Interactome Mapping by Photoaffinity Labeling Reveals Binding Site Hotspots for the NSAIDs. *J. Am. Chem. Soc.*, **140**: 4259.
3. Wang, J., et al. (2015). Mapping sites of aspirin-induced acetylations in live cells by quantitative acid-cleavable activity-based protein profiling (QA-ABPP). *Sci. Rep.*, **5**: 7896.


## Photocleavable Biotin Probes



A major advantage of the photocleavable (PC) linker over all other cleavable linkers is the reagent-free release of the captured biomolecules from streptavidin. This unique property of the photocleavable (PC) linker has promoted its application as a tool for separating, purifying, and identifying desired target biomolecules. PC probes contain a biotin moiety linked to a 'clickable' group through a spacer arm containing a photocleavable moiety. Captured biomolecules can be efficiently photoreleased, typically >90% in 5-25 minutes using an inexpensive, near-UV, low intensity lamp (e.g. 365 nm lamp at 1-5 mW/cm<sup>2</sup>).

### PC Biotin Azide


**CAS:** n/a  
**MW:** 825.37  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Photocleavable biotinylation reagent



**Catalog#**  
**Unit**  
CCT-1119-10 10 mg  
CCT-1119-25 25 mg  
CCT-1119-100 100 mg

### PC Biotin Alkyne


**CAS:** n/a  
**MW:** 780.34  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Photocleavable biotinylation reagent



**Catalog#**  
**Unit**  
CCT-1118-10 10 mg  
CCT-1118-25 25 mg  
CCT-1118-100 100 mg

### PC DBCO Biotin

**CAS:** n/a  
**MW:** 1002.14  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Photocleavable biotinylation reagent




**Catalog#**  
**Unit**  
CCT-1120-10 10 mg  
CCT-1120-25 25 mg  
CCT-1120-100 100 mg

### Selected References:

- Wang, Z., *et al.* (2010). Enrichment and Site Mapping of O-Linked N-Acetylglucosamine by a Combination of Chemical/Enzymatic Tagging, Photochemical Cleavage, and Electron Transfer Dissociation Mass Spectrometry. *Mol. Cell. Proteom.*, **9**: 153.
- Pandor, M., *et al.* (2002). Photochemical Control of the Infectivity of Adenoviral Vectors Using a Novel Photocleavable Biotinylation Reagent. *Chemistry & Biology*, **9**: 567.
- Zhou, G., *et al.* (2010). Photocleavable Peptide-Conjugated Magnetic Beads for Protein Kinase Assays by MALDI-TOF MS. *Bioconjugate Chem.*, **21**: 1917.
- Kim, H., *et al.* (2009). An Azido-Biotin Reagent for Use in the Isolation of Protein Adducts of Lipid-derived Electrophiles by Streptavidin Catch and Photorelease. *Mol. Cell. Proteom.*, **8**: 2080.
- Szychowski, J., *et al.* (2010). Cleavable Biotin Probes for Labeling of Biomolecules via Azide-Alkyne Cycloaddition. *J. Am. Chem. Soc.*, **132**: 18351.


## Dde Biotin Probes



These novel click chemistry probes for enrichment of azide- or alkyne-tagged biomolecules overcome a major drawback of the biotin-streptavidin affinity purification associated with the extraordinary strength of the biotin-streptavidin interaction. These probes contain a biotin moiety linked to a “clickable” group through a spacer arm containing a Dde linker. The Dde moiety is stable to rigorous denaturing wash conditions, such as basic conditions including generally applied buffer systems to which the biological sample may be exposed. At the same time the Dde linker can be quantitatively cleaved under mild aqueous buffered conditions with 2% hydrazine. Finally, the cleaved moiety that remains on the modified peptide minimally changes the peptide mass and generates an additional positive charge, which facilitates peptide sequencing by ETD.

### Dde Biotin Azide


**CAS:** n/a  
**MW:** 695.37  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Cleavable biotinylation reagent



Catalog#	Unit
CCT-1136-1	1 mg
CCT-1136-5	5 mg
CCT-1136-25	25 mg

### Dde Biotin Azide Plus


**CAS:** n/a  
**MW:** 815.98  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Next generation copper-chelating biotin probe for CuAAC..



Catalog#	Unit
CCT-1489-1	1 mg
CCT-1489-5	5 mg
CCT-1489-25	25 mg

### Dde Biotin Picolyl Azide


**CAS:** n/a  
**MW:** 815.98  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Biotinylation reagent with superior kinetics in copper-catalyzed click reactions.



Catalog#	Unit
CCT-1186-1	1 mg
CCT-1186-5	5 mg
CCT-1186-25	25 mg

### Dde Biotin Alkyne

**CAS:** n/a  
**MW:** 650.83  
**Solubility:** DMSO, DMF, THF, DCM  
**Description:** Cleavable biotinylation reagent




Catalog#	Unit
CCT-1137-1	1 mg
CCT-1137-5	5 mg
CCT-1137-25	25 mg

### Selected References:

1. Yang Y., et al. (2013). Cleavable Trifunctional Biotin Reagents for Protein Labeling, Capture, and Release. *Chem. Commun.*, **48**: 5366
2. Matthew E.G., et al. (2016) Comprehensive Mapping of O-GlcNAc Modification Sites Using a Chemically Cleavable Tag. *Mol. Biosyst.* **12**: 1756.
3. Gertsik N., et al. (2017). Mapping the Binding Site of BMS-708163 on  $\gamma$ -Secretase with Cleavable Photoprobes. *Cell Chemical Biology*, **32**: 3.

## Diazo Biotin Probes



Diazobenzene-based biotin probes can be chemoselectively cleaved in mild aqueous buffered conditions with 100 mM sodium dithionite. The diazobenzene linker is stable towards acidic or basic conditions, including generally applied buffer systems to which the biological sample may be exposed.


### Diazo Biotin Azide

CAS: 1339202-33-3

MW: 711.83

Solubility: DMSO, DMF

Description: Cleavable biotinylation reagent



Catalog#

Unit

CCT-1041-1

1 mg

CCT-1041-5

5 mg

CCT-1041-25

25 mg


### Diazo Biotin Alkyne

CAS: n/a

MW: 795.54

Solubility: DMSO, DMF

Description: Cleavable biotinylation reagent



Catalog#

Unit

CCT-1042-1

1 mg

CCT-1042-5

5 mg

CCT-1042-25

25 mg


### Diazo DBCO Biotin

CAS: n/a

MW: 973.15

Solubility: DMSO, DMF,  
DCM, THF

Description: Cleavable  
biotinylation reagent



Catalog#

Unit

CCT-1043-1

1 mg

CCT-1043-5

5 mg

CCT-1043-25

25 mg

### Selected References:


- Yang Y., et al. (2013). Cleavable Trifunctional Biotin Reagents for Protein Labeling, Capture, and Release. *Chem. Commun.*, **48**: 5366
- Yang Y-Y., et al. (2010) Bioorthogonal Chemical Reporters for Monitoring Protein Acetylation. *J. Am. Chem. Soc.* **132**: 3640.
- Rangan K. J., et al. (2010). Rapid visualization and large-scale profiling of bacterial lipoproteins with chemical reporters. *J. Am. Chem. Soc.*, **132**: 10628.

Can't find a product?

Let us know and our team of highly skilled chemists might add it to our catalog

## L-Homopropargylglycine (HPG)


CAS: 98891-36-2  
MW: 163.60  
Solubility: Water, DMSO, DMF



Catalog#	Unit
CCT-1067-25	25 mg
CCT-1067-100	100 mg
CCT-1067-1000	1000 mg

## 4-Azido-L-phenylalanine


CAS: 33173-53-4  
MW: 206.20  
Solubility: Water, DMSO, DMF



Catalog#	Unit
CCT-1406-1G	1 g
CCT-1406-5G	5 g
CCT-1406-25G	25 g

## 5-Ethynyl-2'-deoxyuridine (EdU)


CAS: 61135-33-9  
MW: 252.23  
Solubility: Water, DMSO, DMF



Catalog#	Unit
CCT-1149-25	25 mg
CCT-1149-100	100 mg
CCT-1149-500	500 mg
CCT-1149-1000	1000 mg

## 5-Ethynyl Uridine (5-EU)


CAS: 69075-42-9  
MW: 268.22  
Solubility: Water, DMSO, DMF



Catalog#	Unit
CCT-1261-10	10 mg
CCT-1261-25	25 mg
CCT-1261-100	100 mg
CCT-1261-500	500 mg

## F-ara-EdU


CAS: n/a  
MW: 270.22  
Solubility: Water, DMSO, DMF



Catalog#	Unit
CCT-1403-5	5 mg
CCT-1403-25	25 mg
CCT-1403-100	100 mg
CCT-1403-500	500 mg

## O-propargyl-puromycin (OPP)


CAS: n/a  
MW: 495.54  
Solubility: Water, DMSO, DMF



Catalog#	Unit
CCT-1407-5	5 mg
CCT-1407-25	25 mg
CCT-1407-100	100 mg

**L-Azidohomoalanine (AHA)**

CAS: 942518-29-8  
 MW: 180.59  
 Solubility: Water, DMF, DMSO




Catalog# Unit

CCT-1066-25 25 mg  
 CCT-1066-100 100 mg  
 CCT-1066-1000 1000 mg  
 CCT-1066-5G 5 g

**N-azidoacetylmannosamine tetraacylated (Ac4ManNAz)**

CAS: 361154-30-5  
 MW: 430.37  
 Solubility: DMSO, DMF, MeOH




Catalog# Unit

CCT-1084-5 5 mg  
 CCT-1084-25 25 mg  
 CCT-1084-100 100 mg

**N-(4-pentynoyl)-mannosamine tetraacylated (Ac4ManNAI)**

CAS: 935658-93-8  
 MW: 427.40  
 Solubility: DMSO, DMF, MeOH




Catalog# Unit

CCT-1154-5 5 mg  
 CCT-1154-25 25 mg  
 CCT-1154-100 100 mg

**N-azidoacetylglucosamine tetraacylated (Ac4GlcNAz)**

CAS: 98924-81-3  
 MW: 430.37  
 Solubility: DMSO, DMF, MeOH




Catalog# Unit

CCT-1085-5 5 mg  
 CCT-1085-25 25 mg  
 CCT-1085-100 100 mg

**N-(4-pentynoyl)-glucosamine tetraacylated (Ac4GlcNAI)**

CAS: 1361993-37-4  
 MW: 427.40  
 Solubility: DMSO, DMF, MeOH




Catalog# Unit

CCT-1155-5 5 mg  
 CCT-1155-25 25 mg  
 CCT-1155-100 100 mg

**N-azidoacetylgalactosamine tetraacylated (Ac4GalNAz)**

CAS: 653600-56-7  
 MW: 430.37  
 Solubility: DMSO, DMF, DCM, THF




Catalog# Unit

CCT-1086-5 5 mg  
 CCT-1086-25 25 mg  
 CCT-1086-100 100 mg







**N-(4-pentynoyl)-galactosamine tetraacylated (Ac4GalNAI)**

CAS: 1658458-26-4  
 MW: 427.40  
 Solubility: DMSO, DMF, DCM, THF



Catalog# Unit


CCT-1156-5 5 mg  
 CCT-1156-25 25 mg  
 CCT-1156-100 100 mg

Kdo Azide			Catalog#	Unit
CAS:	1380099-68-2		CCT-1241-10	10 mg
MW:	280.24		CCT-1241-25	25 mg
Solubility:	DMSO, DMF, Water		CCT-1241-100	100 mg
6-azido-6-deoxy-N-acetyl-glucosamine triacylated (Ac3-6AzGlcNAc)			Catalog#	Unit
CAS:	n/a		CCT-1258-5	5 mg
MW:	372.33		CCT-1258-25	25 mg
Solubility:	DMSO, DMF, DCM, THF		CCT-1258-100	100 mg
6-Azide-Trehalose (6-TreAz)			Catalog#	Unit
CAS:	n/a		CCT-1472-5	5 mg
MW:	367.31		CCT-1472-25	25 mg
Solubility:	DMSO, DMF, Water		CCT-1472-100	100 mg
O-Alkyne-Trehalose (O-AlkTMM)			Catalog#	Unit
CAS:	n/a		CCT-1473-5	5 mg
MW:	464.46		CCT-1473-25	25 mg
Solubility:	DMSO, DMF, Water		CCT-1473-100	100 mg
Alkynyl Stearic Acid			Catalog#	Unit
CAS:	34450-18-5		CCT-1166-5	5 mg
MW:	280.45		CCT-1166-25	25 mg
Solubility:	DMSO, DMF, DCM, THF		CCT-1166-100	100 mg
Alkynyl Palmitic Acid			Catalog#	Unit
CAS:	99208-90-9		CCT-1165-5	5 mg
MW:	252.39		CCT-1165-25	25 mg
Solubility:	DMSO, DMF, DCM, THF		CCT-1165-100	100 mg



**Alkynyl Myristic Acid**


CAS: 82909-47-5  
 MW: 244.32  
 Solubility: DMSO, DMF, DCM, THF



Catalog#	Unit
CCT-1164-5	5 mg
CCT-1164-25	25 mg
CCT-1164-100	100 mg

**Azido Myristic Acid**


CAS: 80667-36-3  
 MW: 241.33  
 Solubility: DMSO, DMF, DCM, THF



Catalog#	Unit
CCT-1345-5	5 mg
CCT-1345-25	25 mg
CCT-1345-100	100 mg

**Azido Palmitic Acid**


CAS: 118162-46-2  
 MW: 283.41  
 Solubility: DMSO, DMF, DCM, THF



Catalog#	Unit
CCT-1346-5	5 mg
CCT-1346-25	25 mg
CCT-1346-100	100 mg

**Alkyne Cholesterol**


CAS: 1631985-09-5  
 MW: 396.61  
 Solubility: DMSO, DMF, DCM, THF



Catalog#	Unit
CCT-1409-1	1 mg
CCT-1409-5	5 mg
CCT-1409-25	25 mg

**27-Alkyne Cholesterol**


CAS: 1527467-07-7  
 MW: 396.66  
 Solubility: DMSO, DMF, DCM, THF



Catalog#	Unit
CCT-1410-1	1 mg
CCT-1410-5	5 mg
CCT-1410-25	25 mg

**E-Cholesterol Alkyne**

CAS: n/a  
 MW: 396.66  
 Solubility: DMSO, DMF, DCM, THF



Catalog#	Unit
CCT-1411-1	1 mg
CCT-1411-5	5 mg
CCT-1411-25	25 mg

## Contact Details

---

### Ordering Information

---

Order online at **vectorlabs.com**

Orders may also be placed by email, telephone, or mail.

Please include the following with each order:

- Product name and catalog number
- Unit size and quantity
- Billing and shipping addresses
- Purchase order number
- Name, phone number, address and email address of person placing order

Orders using VISA, Mastercard, or American Express are accepted and processed immediately. Telephone orders over \$1000 may require written confirmation. A confirmation should be boldly marked "Confirming Order. Do Not Duplicate". Duplicate shipments due to incorrectly marked confirming orders cannot be returned for credit. No returned product will be accepted or credited without prior authorization from Vector Laboratories.

Please contact us to discuss discounts for custom or large orders.

### Payment/shipping terms

---

For non-credit card orders, our payment terms are net 30 days from date of invoice, title and risk of loss transfer Ex Works (Incoterms 2010) Seller's location, freight prepaid and added unless shipped on Buyer's account (FedEx, UPS, DHL). Buyers are required to submit a credit application

before credit terms are extended. Orders placed before 3 pm Pacific Time on Monday through Friday (excluding holidays) are usually processed the same day they are received. Unless requested otherwise, all products are shipped 2nd Day Air.



### US Office

---

USA Headquarters:

**Vector Laboratories, Inc.**

**6737 Mowry Avenue**

**Newark, CA 94560**

Telephone Numbers:

Technical Service **(650) 697-3600 x 1**

Customer Service **(650) 697-3600 x 2**

Customer Service: **[vector@vectorlabs.com](mailto:vector@vectorlabs.com)**

Technical Support: **[technical@vectorlabs.com](mailto:technical@vectorlabs.com)**



International Inquiries: **[techintl@vectorlabs.com](mailto:techintl@vectorlabs.com)**

Get in touch with us at

**[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)**

to take the next step in your bioconjugate journey.





Click Chemistry Toolbox

© 2025 Vector Laboratories

